

Quantum Effective Action in Spacetimes with Branes and Boundaries: Diffeomorphism Invariance

A.O.Barvinsky

Theory Department, Lebedev Physics Institute, Leninsky Prospect 53, Moscow 119991, Russia

Abstract

We construct a gauge-fixing procedure in the path integral for gravitational models with branes and boundaries. This procedure incorporates a set of gauge conditions which gauge away effectively decoupled diffeomorphisms acting in the $(d + 1)$ -dimensional bulk and on the d -dimensional brane. The corresponding gauge-fixing factor in the path integral factorizes as a product of the bulk and brane (surface-theory) factors. This factorization underlies a special bulk wavefunction representation of the brane effective action. We develop the semiclassical expansion for this action and explicitly derive it in the one-loop approximation. The one-loop brane effective action can be decomposed into the sum of the gauge-fixed bulk contribution and the contribution of the pseudodifferential operator of the brane-to-brane propagation of quantum gravitational perturbations. The gauge dependence of these contributions is analyzed by the method of Ward identities. By the recently suggested method of the Neumann-Dirichlet reduction the bulk propagator in the semiclassical expansion is converted to the Dirichlet boundary conditions preferable from the calculational viewpoint.

1. Introduction

In this paper we extend the method of quantum effective action in brane models [1] to gravitational systems invariant under general coordinate diffeomorphisms. Specifically, we will be interested in the peculiarities of the gauge-fixing procedure caused by the presence of branes/boundaries. It is well known that branes break the full diffeomorphism invariance and give rise to the dynamical brane bending modes [2] which produce ghost instabilities [3, 4] and generate a low strong-coupling scale [5]. This leads to difficulties in the construction of long-distance modifications of gravity theory that could underly the dark energy phenomenon both at the classical and quantum

level [6]. Quantum effects in gravitational and, in particular, cosmological brane models might also be important for the mechanism of the cosmological acceleration that can be facilitated by the 4D conformal anomaly in the presence of extra dimensions [7]. So here we want to study the gauge-fixing aspects of such problems in the quantum domain.

Peculiarities of the gauge-fixing procedure in brane models follow from the structure of their action

$$S[G_{AB}(X)] = S^B[G_{AB}(X)] + S^b[g_{\alpha\beta}(x)]. \quad (1.1)$$

It contains the bulk and brane parts as functionals of the bulk metric $G_{AB}(X)$ and the induced metric on the brane $g_{\alpha\beta}(x)$. The bulk $(d+1)$ -dimensional and the brane d -dimensional coordinates are labeled respectively by $X = X^A$, $A = 0, 1, \dots, d$, and $x = x^\alpha$, $\alpha = 0, 1, \dots, d-1$, the brane is embedded into the bulk by means of some embedding function $e(x)$,

$$X^A = e^A(x), \quad (1.2)$$

and the induced metric reads in terms of the bulk metric as

$$g_{\alpha\beta}(x) = \partial_\alpha e^A(x) G_{AB}(e(x)) \partial_\beta e^B(x). \quad (1.3)$$

Generically, the bulk part in (1.1) is the Einstein action with the cosmological term

$$S^B[G_{AB}(X)] = \frac{1}{16\pi G} \left\{ \int_{\mathbf{B}} d^{d+1}X G^{1/2} (R(G) - 2\Lambda) - 2 \int_{\mathbf{b}} d^d x g^{1/2} K \right\} \quad (1.4)$$

which includes the integral over the bulk $\mathbf{B}$ and the Gibbons-Hawking integral over the brane $\mathbf{b}$. The latter involves the trace of the brane extrinsic curvature $K = g^{\alpha\beta} K_{\alpha\beta}$ and is necessary for the consistency of the variational procedure for this action.¹

The brane part of (1.4) is given by a covariant d -dimensional integral over $\mathbf{b}$. Depending on the model it contains the brane tension term (in the Randall-Sundrum model [8]), the brane Einstein term (like in the Dvali-Gabadadze-Porrati model [9]), the combination thereof [10] or other covariant structures in brane and bulk curvatures [11]. Both bulk and brane parts of the full action may also contain matter fields which for brevity we do not consider here. Without loss of generality they can be included into the sets of bulk and brane metric coefficients. Also, it is worth noting that Eqs.(1.1)-(1.4) equally well describe the system in the bulk domain $\mathbf{B}$ with the boundary $\mathbf{b} = \partial\mathbf{B}$ and the brane system with the bulk satisfying the Z_2 -symmetry with

¹The extrinsic curvature is defined as a projection onto the brane of the tensor $K_{AB} \equiv \nabla_A n_B$ with the *inward* pointing vector n_B normal to the brane.

respect to the brane $\mathbf{b}$. This formalism also applies in context of Euclidean quantum gravity with a closed compact boundary $\mathbf{b}$.

The bulk and brane parts of the action (1.1) are invariant with respect to diffeomorphisms generated by the bulk $\Xi^A(X)$ and brane $\xi^\alpha(x)$ vector fields. The corresponding transformations have the form

$$\delta^\Xi G_{AB}(X) = \nabla_A \Xi_B(X) + \nabla_B \Xi_A(X), \quad (1.5)$$

$$\delta^\xi g_{\alpha\beta}(x) = D_\alpha \xi_\beta(x) + D_\beta \xi_\alpha(x), \quad (1.6)$$

where ∇_A denotes the covariant derivative in the bulk and D_α is a covariant derivative on the brane. The bulk diffeomorphism (1.5) preserves the bulk action only when the projection of $\Xi^A(X)$ normal to the brane, $\Xi^\perp(X)$, is vanishing. In contrast, the tangential projection of $\Xi^A(X)$, $\Xi^\alpha(X)$, on the brane can be arbitrary. Because the bulk and induced brane metrics are not independent, this projection generates the brane diffeomorphism and coincides with the vector field $\xi^\alpha(x)$. Thus, the boundary conditions for the above gauge transformations read

$$\Xi^\alpha(X)| = \xi^\alpha(x), \quad \Xi^\perp(X)| = 0. \quad (1.7)$$

Here and in what follows the vertical bar indicates that the function in the bulk, $\phi(X)$, is restricted to the brane and labeled by the corresponding low case letter x , $\phi(X)| = \phi(e(x))$, with the aid of the embedding function (1.2).

The construction of the quantum effective action for any brane model should incorporate the gauge-fixing procedure for the diffeomorphism symmetry (1.5)-(1.7). Schematically, this corresponds to introducing the Feynman-DeWitt-Faddeev-Popov gauge-fixing factor in the path integral [12]

$$e^{i\Gamma} = \int DG_{AB}(X) \exp \left(iS[G_{AB}(X)] \right) \times (\text{gauge-fixing}). \quad (1.8)$$

Here integration runs over the metric coefficients in the bulk and on the brane because the metric on the brane is not fixed and is subject to quantum fluctuations. The gauge-fixing factor should gauge away local diffeomorphisms by imposing a certain set of gauge conditions and introducing necessary ghost factors maintaining unitarity and gauge independence of the effective action.

Curiously, despite a well-known form of the gauge algebra (1.5)-(1.7), no exhaustive formulation of gauge-fixing procedure for (1.8) is known in current literature. In particular, the ghost factors require the knowledge of boundary conditions for the corresponding Faddeev-Popov operators acting on the space of gauge parameters $\Xi^A(X)$. In view of (1.7) their tangential components $\Xi^\alpha(X)|$ are arbitrary and should be integrated over, and it would seem that this integration gives rise to the Neumann boundary

conditions for $\Xi^\alpha(X)$. However, this naive prescription turns out to be incorrect. As we show below, the derivation of the boundary conditions for ghosts is less straightforward and much deeper relies on the details of gauging away the transformations (1.5)-(1.7).

In its turn, the preferable choice of gauge conditions in brane models is constrained by additional requirements. One requirement follows from the fact that within the brane concept the fields in the bulk are usually integrated out, so that the physics of the system is effectively probed only by the variables living on the brane.² Thus, a very important aspect of the effective action in brane theory is that it should be a functional of the induced metric $g_{\alpha\beta}(x)$, rather than the full set of metric coefficients $G_{AB}(X)$. Therefore, the preferable gauge conditions should be chosen in such a way that the calculational technique for $\Gamma[g_{\alpha\beta}(x)]$ is manifestly covariant with respect to brane diffeomorphisms (1.6) *separately* from the bulk ones (1.5).

The effective decoupling of brane diffeomorphisms from the bulk diffeomorphisms can be attained by imposing two sets of gauge conditions — brane d -dimensional gauges $\chi^\mu(x) = 0$ and the bulk $(d+1)$ -dimensional gauges $H^A(X) = 0$ which gauge away their corresponding diffeomorphisms. Moreover, they can preserve the manifest covariance of the calculational scheme if they are chosen in the DeWitt background-covariant form [12]. Under the splitting of the brane metric into the background (mean field) and perturbation parts, $g_{\alpha\beta} \rightarrow g_{\alpha\beta} + h_{\alpha\beta}$, this brane gauge reads as

$$\chi^\mu = D^\nu h_{\mu\nu} - \frac{1}{2} D^\mu h, \quad h \equiv g^{\mu\nu} h_{\mu\nu}, \quad (1.9)$$

where the covariant derivatives D_μ are defined with respect to the background metric $g_{\mu\nu}$ which also raises the indices of $h_{\mu\nu}$ and D_μ . The corresponding Faddeev-Popov operator which we denote by $\mathbf{J}_\nu^\mu(D)$ acts on the vector space of $\xi^\nu(x)$. In the lowest order in $h_{\mu\nu}$ it reads as a covariant d -dimensional d'Alembertian modified by the Ricci-curvature potential term

$$\mathbf{J}_\nu^\mu(D) = \square^{(d)} \delta_\nu^\mu + R_\nu^\mu, \quad \square^{(d)} = g^{\mu\nu} D_\mu D_\nu. \quad (1.10)$$

A similar DeWitt bulk gauge condition arises under the splitting of the full bulk metric, $G_{AB} \rightarrow G_{AB} + H_{AB}$, and reads

$$H^A = \nabla^B H_{BC} - \frac{1}{2} \nabla^A H, \quad H \equiv G^{AB} H_{AB}, \quad (1.11)$$

²This means that in the classical action of the path integral (1.8) one must add sources conjugated only to the variables located at the brane, that is to the components of the induced metric $g_{\alpha\beta}(x)$. The Legendre transform with respect to these sources then leads to the effective action as a functional of the mean *induced* metric, $\Gamma[\langle g_{\alpha\beta}(x) \rangle]$. As is well known, in the one-loop approximation this field is called the background, and the whole procedure reduces to the subtraction from $S[G_{AB}(X)]$ in (1.8) the term linear in the deviation of the integration variable from this background — the stationary point of the path integral.

where ∇_A is also defined with respect to the background metric G_{AB} . The corresponding Faddeev-Popov operator, denoted by $Q_B^A(\nabla)$ and acting on the space of $\Xi^B(X)$, is given by

$$Q_B^A(\nabla) = \square^{(d+1)} \delta_B^A + R_B^A, \quad \square^{(d+1)} = \nabla^A \nabla_A. \quad (1.12)$$

This operator is of the second order in derivatives and, as we show below, it should be supplied by the Dirichlet boundary conditions for *all* components of $\Xi^A(X)$. This guarantees the decoupling of the bulk and brane diffeomorphisms even despite their entanglement via the relation (1.7).

Another requirement to the background field formalism for brane theories is the reduction of their Feynman rules to those of the Green's functions with Dirichlet boundary conditions. As discussed in [1] such boundary conditions are computationally much simpler than the original brane matching conditions of the generalized Neumann type. This reduction is possible in view of the duality relations between the Dirichlet and Neumann boundary value problems suggested in [13], and as we show below this reduction also works for gauge-fixed framework. In particular, the decomposition of the quantum effective action in the sum of the bulk and brane-to-brane contributions obtained in [1], which incorporates this Neumann-Dirichlet reduction, works both in the gauge-field and ghost sectors of (1.8).

Below we develop the gauge-fixing procedure of the above type. We derive the gauge-fixing factor in Eq.(1.8), show its bulk-brane factorization property underlying a special bulk wavefunction representation of the brane effective action [14] and build the semiclassical expansion for this action with its Neumann-Dirichlet reduction.

The paper is organized as follows. In Sect.2 we list the main results of this paper. Sect.3 presents the spacetime foliation associated with its brane/boundary and the corresponding canonical formalism which facilitates the derivation of relevant boundary value problems. In Sects. 4 and 5 we derive the gauge-fixing factor of (1.8) and the associated bulk wavefunction representation of the brane effective action. In Sect. 6 we derive the Feynman diagrammatic technique for this action and explicitly present it in the one-loop approximation as a sum of special bulk and brane contributions. They are based on a mixed Dirichlet-Neumann boundary value problem for the graviton propagator. Sect.7 presents the method of Ward identities which allow one to demonstrate for both of these contributions their manifest gauge independence. In Sect. 8 the diagrammatic technique is completely reduced to the propagator with strictly Dirichlet boundary conditions. The concluding section summarizes the obtained results and discusses their possible applications. Three appendices describe the properties of the local measure in the path integral for brane models, the Gaussian integration over the functional space with mixed Dirichlet-Neumann boundary conditions and the variational problem for the Green's function subject to such boundary conditions.

2. Main results

The main results of this paper include the selection of bulk and brane gauge conditions, their corresponding gauge-fixing factor in the path integral (1.8), the one-loop expression for the brane effective action and its Neumann-Dirichlet reduction. In this section we briefly list these results that will be derived in the following sections.

2.1. Bulk and brane gauge conditions

In view of (1.7) a generic diffeomorphism in the bulk can be represented as a composition of two transformations. One transformation is uniquely fixed by a given vector field $\xi^\mu(x)$ on the brane by continuing its d -dimensional diffeomorphism from the brane to the spacetime bulk. Another transformation is a generic $(d+1)$ -dimensional diffeomorphism which reduces to the identity at the boundary, $\Xi^A(X)| = 0$. Correspondingly, the gauge-fixing procedure can be split into two stages. At the first stage special bulk gauge conditions gauge away this $(d+1)$ -dimensional diffeomorphisms vanishing at the brane. At the second stage brane gauge conditions completely fix the residual d -dimensional diffeomorphisms at the boundary. This strategy allows one to decouple the boundary diffeomorphisms from the bulk ones and render the whole formalism manifestly covariant with respect to the both types of diffeomorphisms.

For this purpose the bulk gauge conditions should be special in the sense that they should not overconstrain the relevant $(d+1)$ -dimensional transformations and leave the boundary values $\Xi^\mu(X)| = \xi^\mu(x)$ arbitrary. This means that $\xi^\mu(x)$ play the role of boundary conditions for zero modes of the relevant Faddeev-Popov operator — the transformations which leave the bulk gauge conditions invariant. If we denote these $(d+1)$ -dimensional gauge conditions by

$$H^A(X) = H^A(G_{CD}(X), \partial_B G_{CD}(X)) \quad (2.1)$$

then their gauge transformation serves as the definition of the bulk Faddeev-Popov operator $Q_B^A(\nabla)$

$$\delta^\Xi H^A(X) = Q_B^A(\nabla) \Xi^B(X). \quad (2.2)$$

To fix the bulk diffeomorphisms this operator should be invertible under the Dirichlet boundary conditions $\Xi^A(X)| = 0$, but it must have nontrivial zero modes $\Xi_0^B(X)$, $Q_B^A(\nabla) \Xi_0^B(X) = 0$, subject to inhomogeneous boundary conditions (1.7) “enumerated” by all possible brane vector fields $\xi^\mu(x)$. This is possible when the Faddeev-Popov operator in (2.2) is of the second order in derivatives transversal to the spacetime boundary.

The gauge conditions which generate the Faddeev-Popov operators with such properties can be background-covariant from the bulk $(d+1)$ -dimensional viewpoint and, in particular, given by the DeWitt gauge in the bulk (1.11).

The residual gauge transformations with the parameters ξ^μ are gauged away by imposing the brane gauge conditions $\chi^\mu(x) = 0$. In order to decouple the boundary diffeomorphisms from the bulk ones, one must impose these gauge conditions only upon the brane metric

$$\chi^\mu(x) = \chi^\mu(g_{\alpha\beta}(x), \partial_\mu g_{\alpha\beta}(x)). \quad (2.3)$$

Similarly to (2.2) they generate the brane Faddeev-Popov operator $\mathbf{J}_\nu^\mu(D)$,

$$\delta^\xi \chi^\mu(x) = \mathbf{J}_\nu^\mu(D) \xi^\nu(x), \quad (2.4)$$

which is now determined entirely in terms of quantities induced on the brane. A particular example of such gauge conditions which are background-covariant from the d -dimensional viewpoint is given by the DeWitt gauge conditions (1.9) with the Faddeev-Popov operator (1.10) which is nondegenerate under appropriate boundary conditions at the infinity of the brane (or in view of the closed compact nature of the boundary in Euclidean context when $\square^{(d)}$ becomes a Laplacian).

2.2. The gauge-fixing factor

For the gauge-fixing procedure (2.1)-(2.4) the gauge-fixing factor in (1.8) factorizes into the product of the corresponding bulk and brane factors [14]

$$(\text{gauge-fixing}) = \delta[H] \text{Det}_D Q \times \delta(\chi) \det \mathbf{J}. \quad (2.5)$$

Here $\delta[H]$ and $\delta(\chi)$ denote respectively the $(d+1)$ -dimensional and d -dimensional functional delta-functions,

$$\delta[H] \equiv \prod_{X \in \mathbf{B}} \prod_A \delta(H^A(X)), \quad (2.6)$$

$$\delta(\chi) \equiv \prod_{x \in \mathbf{b}} \prod_\mu \delta(\chi^\mu(x)), \quad (2.7)$$

and $\text{Det}_D Q$ and $\det \mathbf{J}$ are the corresponding ghost functional determinants of the bulk and brane Faddeev-Popov operators defined by (2.2) and (2.4). To distinguish between the functional dimensionalities of these determinants we denote the determinant of the $(d+1)$ -dimensional theory by $\text{Det} \equiv \text{Det}^{(d+1)}$ and that of the d -dimensional theory by $\det \equiv \text{Det}^{(d)}$.

The subscript in $\text{Det}_D Q$ indicates that the functional determinant of the second-order differential operator $Q_B^A(\nabla)$ is taken subject to the *Dirichlet* boundary conditions on the brane. The variational definition of such a determinant in terms of the Dirichlet Green's function of $Q_B^A(\nabla)$ is given below, see Eq.(4.9). In contrast, boundary conditions for $\mathbf{J}(D)$ in $\det \mathbf{J}$ are not important for us, because the brane boundary is either absent, as in Euclidean context with a closed compact boundary of a bulk domain (the boundary of the boundary), or lies at infinity where the boundary conditions are trivial.

One can use the t'Hooft's method of transition from the degenerate (delta-function type) gauges to the gauge-breaking terms both in the bulk sector

$$\delta [H] \Rightarrow (\text{Det } c_{AB})^{1/2} \exp \left(i S_{\text{gb}}^B[G] \right), \quad (2.8)$$

$$S_{\text{gb}}^B[G(X)] = -\frac{1}{2} \int_{\mathbf{B}} dX H^A(X) c_{AB} H^B(X), \quad (2.9)$$

and brane sector

$$\delta (\chi) \Rightarrow (\det c_{\mu\nu})^{1/2} \exp \left(i S_{\text{gb}}^b[g] \right), \quad (2.10)$$

$$S_{\text{gb}}^b[g(x)] = -\frac{1}{2} \int_{\mathbf{b}} dx \chi^\mu(x) c_{\mu\nu} \chi^\nu(x). \quad (2.11)$$

Here c_{AB} and $c_{\mu\nu}$ are respectively the bulk and brane gauge-fixing matrices. We use for them the same notation differing only by the type of indices (AB vs $\mu\nu$), which should not lead to a confusion in what follows.

The factorization of the gauge-fixing factor (2.5) underlies a special bulk wavefunction representation of the brane effective action, see Eqs. (5.2)-(5.3) below.

2.3. One-loop brane effective action

The semiclassical expansion for the brane effective action within the gauge-fixing procedure of the above type has the form

$$\begin{aligned} \Gamma[g_{\alpha\beta}(x)] = & S^B[G_{AB}^0[g(x)]] + S^b[g_{\alpha\beta}(x)] \\ & + \hbar \Gamma_{1\text{-loop}}[g_{\alpha\beta}(x)] + O(\hbar^2). \end{aligned} \quad (2.12)$$

Its tree-level part (the first line of this equation) follows from the classical action (1.1) calculated on the solution of classical equations of motion, $G_{AB}^0[g(x)]$. This solution satisfies the bulk gauge (2.1) and is subject to the boundary condition on the brane —

the induced d -dimensional metric $g_{\alpha\beta}(x)$,

$$\left. \frac{\delta S^{\text{B}}[G]}{\delta G_{AB}(X)} \right|_{G=G^0} = 0, \quad (2.13)$$

$$H^A(G^0(X), \partial G^0(X)) = 0, \quad (2.14)$$

$$G_{\alpha\beta}^0(X)|_{\text{b}} = g_{\alpha\beta}(x). \quad (2.15)$$

The one-loop part of the action is built in terms of the inverse propagator of the theory in the bulk, which is non-degenerate due to the contribution of the bulk gauge-breaking term

$$F = F^{AB,CD}(\nabla) \delta(X, X') = \frac{\delta^2(S^{\text{B}} + S_{\text{gb}}^{\text{B}})}{\delta G_{AB}(X) \delta G_{CD}(X')}. \quad (2.16)$$

The Green's function of this operator, which determines the bulk propagator,

$$G_{\text{DN}}(X, X') = G_{AB,CD}^{\text{DN}}(X, X'), \quad (2.17)$$

satisfies the following problem with the set of mixed boundary conditions of the generalized Dirichlet-Neumann type

$$F(\nabla) G_{\text{DN}}(X, X') = \delta(X, X'), \quad (2.18)$$

$$G_{\alpha\beta,CD}^{\text{DN}}(X, X')|_X = 0, \quad (2.19)$$

$$\vec{H}(\nabla) G_{\text{DN}}(X, X')|_X = 0. \quad (2.20)$$

Here (2.19) means that the induced metric components of the Green's function ($\alpha\beta$ -components among AB -components) vanish on the brane, whereas the rest of the boundary conditions imply the vanishing of the linearized gauge conditions. The differential operator of linearized gauge conditions, acting upon the Green's function in (2.20), $H^{E,AB}(\nabla) G_{AB,CD}^{\text{DN}}(X, X')$, is defined by

$$\vec{H}(\nabla) \delta(X, X') \equiv \vec{H}^{E,AB}(\nabla) \delta(X, X') = \frac{\delta H^E(X)}{\delta G_{AB}(X')}. \quad (2.21)$$

In terms of these quantities the one-loop effective action reads as the sum of the bulk and brane effective actions both including their relevant ghost contributions (corresponding to the factorization of the gauge-fixing factor (2.5))

$$\begin{aligned} i\Gamma_{1\text{-loop}}[g_{\alpha\beta}(x)] = & -\frac{1}{2} \text{Tr}_{\text{DN}} \ln F + \text{Tr}_{\text{D}} \ln Q \Big|_{G=G_0(g)} \\ & -\frac{1}{2} \text{tr} \ln \mathbf{F}^{\text{DN}} + \text{tr} \ln \mathbf{J}. \end{aligned} \quad (2.22)$$

Here Tr_D and Tr_{DN} denote the functional traces of the bulk theory subject to the Dirichlet and mixed Dirichlet-Neumann (cf. (2.18)-(2.20)) boundary conditions, while tr is a functional trace in the boundary d -dimensional theory. $\mathbf{F}^{DN}$ is the brane-to-brane operator

$$\begin{aligned}\mathbf{F}^{DN} &\equiv \mathbf{F}_{DN}^{\alpha\beta, \gamma\delta}(x, x') \\ &= -\left(\vec{W} G_{DN} \overleftarrow{W} \parallel\right)^{\alpha\beta, \gamma\delta}(x, x') + \kappa^{\alpha\beta, \gamma\delta}(D) \delta(x, x').\end{aligned}\quad (2.23)$$

This is a gauge theory generalization of the brane operator introduced in [1]. The first term here implies that the kernel of the Dirichlet-Neumann Green's function is being acted by the operators $W(\nabla)$ upon both arguments in the directions indicated by arrows. The double vertical bar indicates that both points of the kernel are restricted to the brane and labeled by the corresponding low case letters. That is, if the embedding of the boundary/brane in the bulk is denoted by $X = e(x)$, then this explicitly means:

$$\vec{W} G_{DN} \overleftarrow{W} \parallel(x, x') \equiv \vec{W}(\nabla) G_{DN}(X, X') \overleftarrow{W}(\nabla') \Big|_{X=e(x), X'=e(x')}. \quad (2.24)$$

The first order differential operator $W(\nabla)$ is the Wronskian operator

$$\vec{W}(\nabla) = \vec{W}^{AB, CD}(\nabla), \quad (2.25)$$

which is determined for the symmetric second-order differential operator (2.16) by the following Wronskian relation valid for two arbitrary test functions $\phi_{1,2}(X)$

$$\int_B dX \left(\phi_1 \vec{F}(\nabla) \phi_2 - \phi_1 \overleftarrow{F}(\nabla) \phi_2 \right) = - \int_{\partial B} dx \left(\phi_1 \vec{W}(\nabla) \phi_2 - \phi_1 \overleftarrow{W}(\nabla) \phi_2 \right). \quad (2.26)$$

With these definitions the first term of (2.23) is given by the $(\alpha\beta, \gamma\delta)$ -block of the matrix $(\vec{W} G_{DN} \overleftarrow{W} \parallel)^{AB, CD}$. The second term of (2.23) is a contribution of the brane d -dimensional part of the classical action (1.1) and the brane gauge-breaking term (2.11)

$$\kappa^{\alpha\beta, \gamma\delta}(D) \delta(x, x') = \frac{\delta^2(S^b + S_{gb}^b)}{\delta g_{\alpha\beta}(x) \delta g_{\gamma\delta}(x')}. \quad (2.27)$$

The gauge-breaking term here makes the whole operator (2.23) nondegenerate.

2.4. Neumann-Dirichlet reduction

The equation (2.22) generalizes the technique of [1] to brane gravitational models invariant with respect to local diffeomorphisms. The main goal of [1] was a complete reduction of the calculational technique from Neumann-type boundary conditions to

much simpler Dirichlet ones. However, in the algorithm (2.22) this goal is not yet achieved, because of the complexity of boundary conditions for $G_{\text{DN}}(X, X')$ in (2.18)-(2.20). Such a reduction can be done, and in the one-loop approximation it is given by the expression alternative to (2.22)

$$i\Gamma_{1\text{-loop}}[g_{\alpha\beta}(x)] = -\frac{1}{2} \text{Tr}_D \ln F + \text{Tr}_D \ln Q - \frac{1}{2} \text{tr} \ln \mathbf{F}^D + \text{tr} \ln \mathbf{J}. \quad (2.28)$$

Here the functional traces of all bulk operators are calculated subject to Dirichlet conditions, and the brane-to-brane operator $\mathbf{F}^D$ is given by the expression similar to (2.23) but with the Dirichlet Green's function G_D replacing G_{DN}

$$\mathbf{F}_D^{\alpha\beta, \gamma\delta}(x, x') = -(\vec{W} G_D \overleftarrow{W} ||)^{\alpha\beta, \gamma\delta}(x, x') + \kappa^{\alpha\beta, \gamma\delta}(D) \delta(x, x'). \quad (2.29)$$

3. Canonical formalism in the brane foliation of spacetime

The derivation of the above results is much easier in terms of the DeWitt condensed notations [12]. In these notations the bulk metric is labeled by the condensed index $a = (AB, X)$ including both tensor labels and the *bulk coordinates* X

$$G^a = G_{AB}(X), \quad a = (AB, X). \quad (3.1)$$

If we apply the same convention to the definition of the vector field $\Xi^A = \Xi^A(X)$, $A = (A, X)$, then the gauge invariance of the bulk action can be written down in terms of the generators of diffeomorphisms R_A^a

$$\delta^\Xi G^a = R_A^a \Xi^A, \quad (3.2)$$

$$R_A^a \frac{\delta S^B}{\delta G^a} = 0, \quad (3.3)$$

where the contraction of condensed indices implies also integration over the bulk coordinates X . In order to display the differential structure of these generators for various metric components

$$R_C^a = \vec{R}_{AB,C}(\nabla) \delta(X, X'), \quad \vec{R}_{AB,C}(\nabla) = 2G_{(AC} \nabla_{B)}, \quad (3.4)$$

we will need the canonical formalism of the bulk gravitational action, associated with the brane-type foliation of the full spacetime. In this formalism the role of time is played by the extra-dimensional coordinate y . In view of the spacelike nature of y this

variable has nothing to do with a real dynamical evolution, but the local differential properties of the diffeomorphism transformations are very similar to those of the usual canonical formalism in the physical time $t = X^0$.

So we assume that the bulk is foliated by the surfaces of constant y , whose embedding is determined by the embedding functions, $X^A = e^A(x^\alpha, y)$, including the embedding of the brane (1.2), $e^A(x^\alpha) \equiv e^A(x^\alpha, 0)$. Such a foliation determines the vielbein of vectors e_α^A and n^A respectively tangential and normal to the slices, $e_\alpha^A \equiv \partial_\alpha e^A$, $n_A e_\alpha^A = 0$, $G_{AB} n^A n^B = 1$. It also gives rise to the induced metric on the brane and other surfaces of constant y – the generalization of Eq.(1.2), $g_{\alpha\beta}(x, y) = e_\alpha^A(x, y) G_{AB}(e(x, y)) e_\beta^B(x, y)$.

This foliation generates the lapse and shift functions defined as normal and tangential projections of the local “velocity” vector $\partial_y e^A(x, y) \equiv \dot{e}^A$ with which the bulk slice evolves in “time” y

$$N^\perp = n_A \partial_y e^A(x, y), \quad N^\alpha = g^{\alpha\beta} N_\beta, \quad N_\beta = e_\beta^B G_{BA} \partial_y e^A(x, y). \quad (3.5)$$

In fact these functions are equivalent to G_{Ay} -components of the full metric, so that $G_{AB}(X)$ can be parameterized in terms of $g_{\alpha\beta}$ and $N^A = (N^\alpha, N^\perp)$. Then the bulk Einstein action (1.4) has a well-known ADM form in terms of the extrinsic curvature $K_{\alpha\beta}$ of constant y slices and their scalar curvature $R^{(d)}(g)$

$$S^B[G] = \frac{1}{16\pi G} \int_{y \geq 0} dy \int dx g^{1/2} N^\perp (K^2 - K_{\alpha\beta}^2 + R^{(d)}(g) - 2\Lambda), \quad (3.6)$$

$$K_{\alpha\beta} = \frac{1}{2N^\perp} (\partial_y g_{\alpha\beta} - D_\alpha N_\beta - D_\beta N_\alpha). \quad (3.7)$$

It is important that only the “velocities” of $g_{\alpha\beta}$,

$$\partial_y g_{\alpha\beta} \equiv \dot{g}_{\alpha\beta}, \quad (3.8)$$

enter the Lagrangian, while the lapse and shift functions N^A are not dynamical and serve as Lagrange multipliers in the y -time canonical formalism of the action (3.6). In spacetime condensed notations of (3.1) we will denote the induced metric of y -slices by g^i and this decomposition will look like

$$G^a = (g^i, N^A), \quad a = (i, A), \quad i = (\alpha\beta, X), \quad A = (\alpha, \perp, X), \\ g^i = g_{\alpha\beta}(X), \quad N^A = (N^\alpha(X), N^\perp(X)). \quad (3.9)$$

In what follows we will also need *canonical* condensed notations in which the indices (a, i, A) include together with discrete labels only the *brane coordinates* x , and the contraction of these indices implies the integration only over x .

$$G^a(y) = (g^i(y), N^A(y)), \quad i = (\alpha\beta, x), \quad A = (\alpha, \perp, x), \\ g^i(y) = g_{\alpha\beta}(x, y), \quad N^A(y) = (N^\alpha(x, y), N^\perp(x, y)) \quad (3.10)$$

In these notations the generators of Eq.(3.2), R_A^a , form delta-function type kernels in the variable y with two entries $a \rightarrow (a, y)$, $A \rightarrow (A, y')$,

$$R_A^a = R_A^a(\partial_y) \delta(y - y'). \quad (3.11)$$

Various components of $R_A^a(\partial_y)$ are either the ultralocal (multiplication) or differential operators acting on the first argument of the delta function.

For sake of brevity, when using the condensed notations of the canonical or space-time nature we will not introduce special labels to distinguish between them. As a rule, when the y -argument is explicitly written we imply that the corresponding condensed indices are canonical, i.e. they contain only discrete labels and brane coordinates x , and their contraction does not involve implicit y -integration. For example, the left-hand side of (3.3) can be written down in the form

$$R_A^a \frac{\delta S^B}{\delta G^a} = \overrightarrow{R_A^a}(\partial_y) \frac{\delta S}{\delta G^a(y)}, \quad A \rightarrow (A, y), \quad (3.12)$$

where the integration over y (implicit in the contraction of the *spacetime condensed* index a) removed the delta function contained in R_A^a and the result boiled down to the action of the differential operator $R_A^a(\partial_y)$ on $\delta S/\delta G^a(y)$. This operator obviously differs from that of Eqs.(3.2) and (3.11) by the functional transposition – integration by parts, because in contrast to (3.2) it acts on the test function with respect to the upper index a . This fact is indicated by the order of operator indices reversed relative to Eq. (3.11).

Another distinction between these two types of condensed notations concerns functional derivatives. We shall always reserve the functional variational notation $\delta/\delta G^a \equiv \delta/\delta G^a(y)$ for the variational derivative with respect to the functions of y , while the variational derivative with respect to the functions of brane coordinates will be denoted by partial derivatives. For example, $\delta/\delta G^a \equiv \delta/\delta G_{AB}(X)$ vs $\partial/\partial g^i \equiv \delta/\delta g_{\alpha\beta}(x)$.

In these notations the action has the form

$$S^B[G] = \int dy L^B(g, \dot{g}, N) \quad (3.13)$$

where the ADM Lagrangian is displayed explicitly depending on $\dot{g}^i = \dot{G}^i$, and all x -derivatives are implicit in condensed canonical notations. With the definition of the momenta conjugated to g^i

$$p_i = p_i^0(g, \dot{g}, N) \equiv \frac{\partial L^B(g, \dot{g}, N)}{\partial \dot{g}^i} \quad (3.14)$$

the bulk action can be rewritten in the canonical form

$$S^B[g, p, N] = \int dy (p_i \dot{g}^i - N^A \mathbf{H}_A(g, p)), \quad (3.15)$$

$$S^B[g, p, N] \Big|_{p=p^0(g, \dot{g}, N)} = S^B[G], \quad (3.16)$$

where $\mathbf{H}_A(g, p) = (\mathbf{H}_\alpha(g, p), \mathbf{H}_\perp(g, p))$ is a set of momentum and Hamiltonian constraints. Similarly to the ADM formalism in the physical time, these constraints as functions on the phase space of (g^i, p_i) satisfy the Poisson bracket relations

$$\{\mathbf{H}_A, \mathbf{H}_B\} = U_{AB}^C \mathbf{H}_C \quad (3.17)$$

with structure functions $U_{AB}^C = U_{AB}^C(g)$, signifying that these constraints belong to the first class according to the Dirac classification. As a consequence of (3.16) they also comprise the $(\perp A)$ -projections of the Einstein equations in the bulk,

$$\mathbf{H}_A(g, p) \Big|_{p=p^0(g, \dot{g}, N)} = -\frac{\delta S^B[g, N]}{\delta N^A}. \quad (3.18)$$

Due to the constraint algebra the canonical action (3.15) is invariant under the gauge transformations with local (arbitrary time and space dependent) parameters $\Xi^A(X)$ satisfying $|\Xi^\perp(X)| = 0$.³ These transformations are canonical and, therefore, ultralocal in y for phase space variables, but involve the y -derivative of $\Xi^A(X)$ for Lagrange multipliers [15]

$$\delta^\Xi g^i = \{g^i, \mathbf{H}_A\} \Xi^A, \quad \delta^\Xi p_i = \{p_i, \mathbf{H}_A\} \Xi^A, \quad (3.19)$$

$$\delta^\Xi N^A = \dot{\Xi}^A - U_{BC}^A N^B \Xi^C. \quad (3.20)$$

In view of the relation (3.16) between the canonical and Lagrangian formalisms various components of R_A^a follow from the transformations (3.19)-(3.20) [15]

$$R_A^i = \delta(y - y') \frac{\partial \mathbf{H}_A}{\partial p_i} \Big|_{p=p^0(g, \dot{g}, N)}, \quad (3.21)$$

$$R_A^B = \vec{R}_A^B(\partial_y) \delta(y - y') = (\delta_A^B \partial_y - U_{CA}^B N^C) \delta(y - y'). \quad (3.22)$$

The distinguished role of the Lagrange multipliers manifests itself in the fact that only the $a = B$ component of (3.11) forms the first order differential operator while the $a = i$ components are ultralocal in y .

Sometimes a composition of differential operators results in an ultralocal operator. Here is one important example that follows from the transformation property of the momentum. On the one hand it is given by the canonical transformation

$$\delta^\Xi p_i = -\frac{\partial \mathbf{H}_A}{\partial g^i} \Xi^A. \quad (3.23)$$

³Tangential to the brane components $|\Xi^\mu| = \xi^\mu$ should not necessarily vanish because these diffeomorphisms do not shift the boundary. In the canonical formalism this property follows from the linearity of the momentum constraints H_μ in p , due to which relevant surface terms identically vanish for arbitrary $|\Xi^\mu|$.

On the other hand it can be obtained by the gauge transformation of the Lagrangian expression for the momentum (3.14). The metric variation of the latter has the form of the differential operator acting on $\delta G^a(y) = (\delta g^i(y), \delta N^A(y))$,

$$\delta_G \frac{\partial L^B(g, \dot{g}, N)}{\partial \dot{g}^i} \equiv W_{ia}^S(\partial) \delta G^a(y). \quad (3.24)$$

In essence $W_{ia}^S(\partial)$ here is a part of the Wronskian operator (2.25)) associated with the part of the operator (2.16), $F_{ab} = F^{AB, CD}(\nabla) \delta(X, X')$, without the gauge-breaking term. Substituting the gauge variation of the metric one therefore has

$$\delta^\Xi p_i^0(g, \dot{g}, N) = \vec{W}_{ia}^S \delta^\Xi G^a(y) = \vec{W}_{ia}^S \vec{R}_A^a \Xi^A \quad (3.25)$$

and comparing (3.23) and (3.25) finds that the composition of the two first-order differential operators is ultralocal in y (and contains at most the derivatives with respect to brane coordinates)⁴

$$\vec{W}_{ia}^S(\partial) \vec{R}_A^a(\partial) = - \left. \frac{\partial \mathbf{H}_A}{\partial g^i} \right|_{p=p^0(g, \dot{g}, N)} \quad (3.26)$$

The final comment of this section concerns the diffeomorphisms tangential to the brane. In the set of $R_A^i = (R_\mu^i, R_\perp^i)$ their generators $R_\mu^i = R_\mu^i(g)$ are independent of $\dot{g}^i$ and correspond to the momentum constraints linear in p ,

$$\mathbf{H}_\mu(g, p) = R_\mu^i p_i. \quad (3.27)$$

By construction they leave the brane part of the action invariant

$$R_\mu^i \frac{\partial S^b}{\partial g^i} = 0, \quad (3.28)$$

which is the brane, d -dimensional, analogue of (3.3).

4. Gauge-fixing procedure

The relativistic nature of the bulk gauge conditions (2.1) which in condensed notations we denote by

$$H^A = H^A(G^a(y), \dot{G}^a(y)) \quad (4.1)$$

⁴The equivalence of the canonical and Lagrangian gauge transformations of momenta, $(\delta^\Xi p)|_{p^0(g, \dot{g}, N)} = \delta^\Xi(p^0(g, \dot{g}, N))$, holds as it follows from (3.16) only up to terms proportional to equations of motion [15]. In what follows the bulk background always satisfies on-shell condition which justifies the identification of canonical and Lagrangian versions of gauge transformations.

implies that they necessarily depend on the “velocities” of the Lagrange multipliers $\dot{N}^A = \partial_y N^A$, so that the following matrix is nondegenerate

$$a_B^A = -\frac{\partial H^A}{\partial \dot{N}^B}, \quad \det a_B^A \neq 0. \quad (4.2)$$

As a result the corresponding Faddeev-Popov operator in view of the derivative nature of the transformations (3.20) for N^A is of the second order in ∂_y . This follows from the definition of this operator

$$Q_B^A = \frac{\delta H^A}{\delta G^a} R_B^a = \vec{H}_a^A(\partial) \vec{R}_B^a(\partial) \delta(y - y'), \quad (4.3)$$

$$\frac{\delta H^A}{\delta G^a} \equiv H_a^A = \vec{H}_a^A(\partial) \delta(y - y'), \quad (4.4)$$

where H_a^A is the matrix of linearized gauge conditions (2.21)). This functional matrix in relativistic gauges is the first-order differential operator, $H_C^A(\partial_y) = -a_C^A \partial_y + \dots$, whence in view of (3.22) $Q_B^A(\partial_y) = -a_B^A \partial_y^2 + \dots$. Such a second-order operator admits zero modes parameterized by their boundary conditions, corresponding to the residual transformations discussed in Introduction and in Sect.2.

To fix these residual transformations we impose gauge conditions on the brane metric (2.3), $\chi^\mu(g) = 0$. They force these boundary conditions to vanish, provided the brane Faddeev-Popov operator, which in condensed canonical notations reads as

$$\mathbf{J}_\nu^\mu = \frac{\partial \chi^\mu}{\partial g^i} R_\mu^i, \quad (4.5)$$

is nondegenerate. As discussed in Introduction, the brane gauge conditions can be chosen background-covariant from the d -dimensional viewpoint, as is the case of the brane DeWitt gauge (1.9). However, they are imposed only on $g^i = g_{\alpha\beta}(x)$ and do not involve $\dot{g}^i$, so that they can be considered unitary from the viewpoint of the y -time canonical formalism.

Thus, the overall gauge-fixing factor in the path integral (1.8) takes the form

$$(\text{gauge-fixing}) = \delta [H^A(G)] \delta(\chi^\mu(g)) M_{H, \chi}[G], \quad (4.6)$$

where we use different condensed notations for the functional delta-functions in the bulk (2.6) and on the brane (2.7). The measure factor $M_{H, \chi}[G]$ is determined according to the standard Faddeev-Popov procedure by the following functional integral over the full diffeomorphism group

$$(M_{\chi, H}[G])^{-1} = \int_{\Xi^\perp| = 0} D\Xi \delta [H^A(G_\Xi)] \delta(\chi^\mu(g_\Xi)). \quad (4.7)$$

To calculate it we write the infinitesimally transformed bulk and brane metrics as $G_{\Xi}^a = G^a + R_A^a \Xi^A + O[\Xi^2]$ and $g_{\xi}^i = g^i + R_{\mu}^i \xi^{\mu} + O[\xi^2]$, and then decompose in (4.7) the integration over $\Xi^A(X)$ into the integration over the bulk field with fixed boundary values $\Xi^A(X)| = (0, \xi^{\mu}(x))$ and the subsequent integration over $\xi^{\mu}(x)$

$$\begin{aligned} \int_{\Xi^{\perp}|=0} D\Xi \delta[H^A(G_{\Xi})] \delta(\chi^{\mu}(g_{\xi})) &= \int d\xi \delta(\mathbf{J}_{\nu}^{\mu} \xi^{\nu}) \int_{\substack{\Xi^{\perp}|=0 \\ \Xi^{\mu}|=\xi^{\mu}}} D\Xi \delta[Q_B^A \Xi^B] \\ &= (\det \mathbf{J}_{\nu}^{\mu})^{-1} \int_{\Xi^A|=0} D\Xi \delta[Q_B^A \Xi^B] = (\det \mathbf{J}_{\nu}^{\mu})^{-1} (\text{Det}_{\text{D}} Q_B^A)^{-1}. \end{aligned} \quad (4.8)$$

The result is structurally very simple – it factorizes into the product of functional determinants of brane and bulk Faddeev-Popov operators, and the latter, $\text{Det}_{\text{D}} Q_B^A$, is calculated subject to Dirichlet conditions for *all* components of $\Xi^A|$. This directly follows from the gauge-fixing procedure on the brane.

The Dirichlet type functional determinant is determined by the variational formula

$$\delta \ln \text{Det}_{\text{D}} Q_B^A = Q_{\text{D}A}^{-1B} \overleftarrow{\delta} Q_B^A \equiv \int dy Q_{\text{D}A}^{-1B}(y', y) \overleftarrow{\delta} Q_B^A(\partial_y) \Big|_{y'=y}, \quad (4.9)$$

where $\overleftarrow{\delta} Q_B^A$ denotes the variation of the operator under generic change of its coefficients and $Q_{\text{D}A}^{-1B}$ is the Dirichlet Green's function of the ghost operator, defined by

$$\begin{aligned} Q_B^C(\partial_y) Q_{\text{D}A}^{-1B}(y, y') &= \delta_A^C \delta(y - y'), \\ Q_{\text{D}A}^{-1B}(y, y') \Big|_y &= 0. \end{aligned} \quad (4.10)$$

Thus, this derivation yields one of the main results of this paper (2.5) and confirms boundary conditions for the ghost propagator in the bulk.

As a result of transition (2.8)-(2.11) the quantum effective action takes the form

$$e^{i\Gamma} = \int DG \mu[G] \exp \left(i S_{\text{gf}}[G] \right) \det \mathbf{J}_{\nu}^{\mu} \text{Det}_{\text{D}} Q_B^A, \quad (4.11)$$

$$S_{\text{gf}}[G] = S^{\text{B}}[G] + S_{\text{gb}}^{\text{B}}[G] + S^{\text{b}}[g] + S_{\text{gb}}^{\text{b}}[g], \quad (4.12)$$

with the full gauge-fixed action $S_{\text{gf}}[G]$ including the bulk and brane gauge-breaking terms which read in canonical condensed notations as

$$S_{\text{gb}}^{\text{B}}[G] = -\frac{1}{2} \int dy H^A(G, \dot{G}) c_{AB} H^B(G, \dot{G}), \quad (4.13)$$

$$S_{\text{gb}}^{\text{b}}(g) = -\frac{1}{2} \chi^{\mu}(g) c_{\mu\nu} \chi^{\nu}(g). \quad (4.14)$$

The last comment of this section concerns the local measure denoted in (4.11) by $\mu[G]$. Its contribution is not very important for practical purposes, because its function solely consists in the cancellation of strongest power divergences of the path integral. Nevertheless, for completeness we briefly discuss it here.

The local measure is determined by another type of bulk and brane foliation – the one with the slices of constant physical time $t = X^0$. This foliation determines the chronological ordering in the unitary evolution in physical (not fictitious) time. It gives rise to the local measure as a contribution of the gaussian path integral over the physical momenta conjugated to temporal velocities $\partial_0 G$ (rather than to $\partial_y G$). As shown in Appendix A, when the brane action has the kinetic term⁵, the full local measure factorizes into the product of the bulk and brane measures

$$\mu[G] = \mu_B[G] \mu_b(g). \quad (4.15)$$

The bulk measure looks as follows. In the foliation of the bulk by spacelike slices of constant $X^0 = t$, $X = (t, \mathbf{x}, y)$ (where $\mathbf{x}$ denotes spatial coordinates among brane coordinates x), the bulk metric can be decomposed into the corresponding spacelike metric $Q(t, \mathbf{x}, y)$ and bulk lapse and shift functions $\mathcal{N}(t, \mathbf{x}, y) \sim G_{At}(X)$, $G_{AB}(X) \sim (Q(t, \mathbf{x}, y), \mathcal{N}(t, \mathbf{x}, y))$. The bulk measure then reads as

$$\mu_B[G] = \prod_{X, X \notin \mathbf{b}} \left[\det \frac{\partial^2 L_B(Q, \partial_t Q, \mathcal{N})}{\partial(\partial_t Q) \partial(\partial_t Q)} \det c_{AB} \right]^{1/2}. \quad (4.16)$$

It also absorbs the determinant of the gauge-fixing matrix c_{AB} which is generated by the transition (2.8)-(2.11) to nondegenerate gauges and which we also consider ultralocal in X .

Similarly, in the construction of the brane measure one has a foliation of the brane by spacelike $(d-1)$ -dimensional surfaces $x = (t, \mathbf{x})$, which leads to $((d-1)+1)$ -decomposition $g(x) = (q(t, \mathbf{x}), n(t, \mathbf{x}))$ into the space metric $q(t, \mathbf{x})$ and brane lapse and shift functions $n(t, \mathbf{x})$. The brane Lagrangian then depends on velocities of only the q -metric coefficients, $L_b(g, \partial g) = L_b(q, \partial_t q, n)$, and the brane measure takes the form

$$\mu_b(g) = \prod_x \left[\det \frac{\partial^2 L_b}{\partial(\partial_t q) \partial(\partial_t q)} \det c_{\mu\nu} \right]^{1/2}. \quad (4.17)$$

⁵Which is not always the case – in the Randall-Sundrum model, for example, the brane part of the action is kinetically inert, because it contains only the brane tension.

5. Bulk wavefunction representation of the brane effective action

As a next step we decompose the full integration in (4.11) into the integration over the bulk metric $G_{AB}(X)$ subject to fixed induced metric on the brane $g_{\alpha\beta}(x) = G_{\alpha\beta}(X)|$ and the subsequent integration over d -dimensional $g_{\alpha\beta}(x)$,

$$\int DG_{AB}(X) = \int dg_{\alpha\beta}(x) \int_{G_{\alpha\beta}|=g_{\alpha\beta}(x)} DG_{AB}(X) \equiv \int dg \int_{G^i|=g^i} DG. \quad (5.1)$$

The result looks as the Feynman-DeWitt-Faddeev-Popov functional integral [14]

$$e^{i\Gamma} = \int dg \mu_b[g] \exp i(S^b(g) + S_{\text{gb}}^b(g)) \det \mathbf{J}_\nu^\mu \Psi_B(g) \quad (5.2)$$

for the purely d -dimensional system with the brane action $S^b[g_{\alpha\beta}]$ but with the insertion of the functional $\Psi_B(g)$ which we will call a wavefunction of the bulk spacetime

$$\Psi_B(g) = \int_{G^i|=g^i} DG \mu_B[G] \exp i(S^B[G] + S_{\text{gb}}^B[G]) \text{Det}_D Q_B^A. \quad (5.3)$$

This function is well known from quantum cosmology as a path-integral representation of the solution of the Wheeler-DeWitt equations – quantum Dirac constraints on a quantum state in the canonical quantization of gravity. At various levels of rigor and in various contexts it was built in a path-integral form in [16, 17, 18, 19, 20]. In particular, in the $d = 3$ context it was obtained in [18] where the initial and final spacelike Cauchy surfaces played the role of branes with specified 3-metrics. The metrics served as functional arguments of the two-point kernel of the unitary evolution interpolating between these two 3-surfaces.

In (5.3) the role of the boundary is played by timelike brane(s), and the canonical formalism in real time is generalized to the case of the “evolution” in the direction transversal to the brane. Certainly, no sensible unitarity or causality can be ascribed to this evolution. Apparently this construction can be generalized to a purely Euclidean case when the full boundary can have various topological and connectedness properties.

The bulk wavefunction (5.3) has the same properties as the cosmological wavefunction of [18, 20]. First of all, it is independent of the choice of the bulk gauge H^A ,

$$\delta_H \Psi_B(g) = 0, \quad (5.4)$$

though H^A explicitly enters its construction. This is a typical property of the Faddeev-Popov functional integral on shell (or in the absence of sources located in the bulk — the only source of $\Psi_B(g)$ is the brane metric $g = g_{\alpha\beta}(x)$). A formal proof is based on the change of the integration variable in (5.3)

$$G^a \rightarrow G_{\Xi}^a = G^a + R_A^a \Xi^A, \quad \Xi^A = Q_B^{-1A} \delta H^B, \quad (5.5)$$

simulating the change of gauge conditions $H^A \rightarrow H^A + \delta H^A$. Due to Ward identities for ghost and gauge Green's functions and gauge invariance of the bulk action, this transformation is identical and leads to (5.4). Important point of the transform (5.5) is that the Green's function of the Faddeev-Popov operator Q_B^{-1A} has Dirichlet boundary conditions, and therefore $\Xi^A| = 0$, so that this transform does not shift g^i on the brane (as it should). It only shifts lapse and shift functions $N^C|$ because of the y -derivative in $R_A^C(\partial_y)$ acting on $\Xi^A(y)$, but this is not dangerous because the Lagrange multipliers $N^C|$ are integrated over at the boundary, rather than being fixed like $G^i = g^i$.⁶

Important consequence of this integration over $N^C|$ is that $\Psi_B(g)$ satisfies the analogue of the Wheeler-DeWitt equations

$$\hat{H}_A(g, \partial/i\partial g) \Psi_B(g) = 0, \quad (5.6)$$

where $\hat{H}_A(g, \partial/i\partial g)$ are the operators of quantum Dirac (momentum and Hamiltonian) constraints. They follow from their classical counterparts by some nontrivial operator realization which is formally known only in the linear in $\hbar$ approximation [21].⁷

The momentum components of (5.7), $A = \mu$, describe the gauge invariance of $\Psi_B(g)$ under d -dimensional diffeomorphisms. In condensed notations this reads as

$$R_{\mu}^i \frac{\partial \Psi_B}{\partial g^i} = 0. \quad (5.7)$$

This property can also be independently proven by the transformation of integration variable similar to (5.5) but with $\Xi^{\mu}| = \xi^{\mu} \neq 0$ — the diffeomorphism tangential to

⁶This is a point of departure from [5] where the effective action is a functional of $N^C|$ and, therefore, depends on the choice of the gauge in the bulk.

⁷It is not obvious, though, that the realization of [21] for the temporal constraints would work for brane constraints, because the latter govern an artificial evolution in the y -time which is not related to unitarity and causality. In particular, the choice of the local measure in (5.3) is not related to y -foliation of the bulk and does not yield the scalar density nature of $\Psi_B(g)$ under the reparameterizations of g . This choice is tightly related to the operator realization of quantum Dirac constraints. For the temporal constraints of [21] the choice of measure guarantees that the cosmological wavefunction is a scalar density of 1/2-weight, and the operators $\hat{H}_A(g, \partial/i\partial g)$ act covariantly on this density. No such properties hold for $\Psi_B(g)$ — a scalar in g rather than a density, see Appendix A.

the brane [19]. This property in its turn guarantees the χ -gauge independence of the *on-shell* brane action Γ ,

$$\delta_\chi \Gamma = 0. \quad (5.8)$$

Indeed, the d -dimensional analogue of the change of integration variables (5.5) in the path integral (5.2) gives this property in virtue of gauge invariance of $S^b(g)$ and $\Psi_B(g)$ and the Ward identities for the brane Faddeev-Popov Green's function.

6. Semiclassical expansion

6.1. Wavefunction of the bulk

Semiclassical expansion for $\Psi_B(g)$ is based on the stationary point of the gauge-fixed action in the bulk. As a result of integration by parts, the variation of the action includes the bulk and boundary terms

$$\begin{aligned} \delta (S^B[G] + S_{\text{gb}}^B[G]) &= \int_B dX \left(\frac{\delta S^B}{\delta G^a(X)} - \frac{\vec{\delta H}^D}{\delta G^a(X)} c_{DB} H^B \right) \delta G^a(X) \\ &\quad + \left(\frac{\partial L_B}{\partial \dot{G}^a} - \frac{\partial H^D}{\partial \dot{G}^a} c_{DB} H^B \right) \delta G^a \Big| \end{aligned} \quad (6.1)$$

which separately should be equated to zero. Now, take into account that for fixed $G^i| = g^i$ the boundary variation $\delta G^i| = 0$ and note that L_B is independent of $\dot{N}^A$ and the matrices $\partial H^D / \partial \dot{N}^A = -a_A^D$ (cf. Eq.(4.2)) and c_{DB} are invertible. Therefore, the stationarity conditions reduce to

$$\frac{\delta S^B}{\delta G^a(X)} - \frac{\delta H^D}{\delta G^a(X)} c_{DB} H^B = 0, \quad (6.2)$$

$$H^B| = 0. \quad (6.3)$$

Functionally contracting (6.2) with R_A^a and using gauge invariance of the bulk action (3.3) we get for the gauge conditions the homogeneous equation with the second-order Faddeev-Popov operator

$$Q_A^D(\partial) c_{DB} H^B(X) = 0. \quad (6.4)$$

Since in view of (6.3) they have Dirichlet boundary conditions, the solution is identically zero everywhere in the bulk. Thus, eventually the stationary point of the bulk action

$$G_0^a(g) \equiv G_{AB}^0(X)[g_{\alpha\beta}(x)] \quad (6.5)$$

is a solution of classical equations of motion in the bulk gauge with fixed brane metric at the boundary, as advocated in Eqs.(2.13)-(2.15) of Sect.2.

The quadratic part of the action on this background – the variation of (6.1) – reads as

$$\frac{1}{2} \delta^2 (S^B[G] + S_{\text{gb}}^B[g]) = \frac{1}{2} \delta G^a \left(\vec{F}_{ab} \delta G^b \right) + \frac{1}{2} \delta G^a \vec{W}_{ab}(\partial) \delta G^b, \quad (6.6)$$

where $\vec{F}_{ab}$ is the condensed notation for the operator (2.16)

$$F_{ab} = S_{ab} - H_a^A c_{AB} H_b^B, \quad (6.7)$$

$$S_{ab} \equiv \vec{S}_{ab}(\partial_y) \delta(y - y') = \frac{\delta^2 S^B[G]}{\delta G^a(y) \delta G^b(y')} \quad (6.8)$$

and $\vec{W}_{ab}(\partial)$ is the corresponding Wronskian operator (2.25) defined by the variational relation

$$\begin{aligned} \vec{W}_{ab}(\partial) \delta G^b(y) &= \delta \left(\frac{\partial L^B}{\partial \dot{G}^a} - \frac{\partial H^A}{\partial \dot{G}^a} c_{AB} H^B \right)_{H^A=0} \\ &= \left(\vec{W}_{ab}^S(\partial) - \frac{\partial H^A}{\partial \dot{G}^a} c_{AB} \vec{H}_b^B(\partial) \right) \delta G^b(y). \end{aligned} \quad (6.9)$$

Together with the Wronskian operator of the invariant action $W_{ab}^S(\partial)$, see Eq. (3.24), it includes the contribution of the gauge-breaking term. $\vec{W}_{ab}(\partial)$ participates in the Wronskian relation (2.25) and also satisfies an additional relation that can be obtained by a single integration by parts of the derivatives in $\vec{F}_{ab}(\partial)$

$$\phi_1^a \vec{F}_{ab} \phi_2^b = \phi_1^a \overleftrightarrow{F}_{ab} \phi_2^b - \phi_1^a \vec{W}_{ab}(\partial) \phi_2^b \quad (6.10)$$

Here $\overleftrightarrow{F}_{ab}$ implies that the derivatives in $\vec{F}_{ab}(\partial)$ are integrated by parts one time to form the bilinear combinations of the first-order derivatives of $\phi_{1,2}$ (for $\phi_2 = \phi_1$ this is just the Lagrangian quadratic in ϕ and $\partial\phi$). With this relation the quadratic form of the action (6.6) takes the form

$$\frac{1}{2} \delta^2 (S^B[G] + S_{\text{gb}}^B[g]) = \frac{1}{2} \delta G^a \overleftrightarrow{F}_{ab} \delta G^b. \quad (6.11)$$

The one-loop contribution of this form to the path integral (5.3) is the gaussian functional integral over the perturbations $\delta G^a = \delta G_{AB}(X)$ subject to Dirichlet boundary conditions for the brane metric components $\delta G^i| = 0$, while $\delta N^A|$ are integrated over in the infinite limits. As shown in Appendix B this integral yields the functional

determinant of F_{ab} subject to mixed Dirichlet-Neumann boundary conditions, and the one-loop wavefunction of the bulk has the final form

$$\Psi_B(g) = \frac{\mu_B \text{Det}_D Q_B^A}{(\text{Det}_{\text{DN}} F_{ab})^{1/2}} \exp \left(i S^B[G] \right) \Big|_{G=G_0(g)}. \quad (6.12)$$

Here the Dirichlet-Neumann functional determinant $\text{Det}_{\text{DN}} F_{ab}$ is determined by the variational relation

$$\delta \ln \text{Det}_{\text{DN}} F_{ab} \equiv \delta \text{Tr}_{\text{DN}} \ln F_{ab} = \overset{\leftrightarrow}{\delta} F_{ab} G_{\text{DN}}^{ba}. \quad (6.13)$$

The notation $\overset{\leftrightarrow}{\delta} F_{ab}$ means arbitrary variations of the coefficients of the operator, and, similarly to Eq.(6.10), the double arrow implies symmetric action of two first-order derivatives of $\delta F_{ab} = \delta F^{AB,CD}(\nabla)$ on both arguments of the Green's function $G_{\text{DN}}^{ba} = G_{CD,AB}^{\text{DN}}(X, X')$ (before taking the coincidence limit $X' = X$ implicit in the functional trace operation).

The most peculiar element of the variational definition (6.13) is G_{DN}^{ba} – the Green's function of the operator $F_{ab}(\partial)$ subject to the mixed set of Dirichlet and generalized Neumann boundary conditions (2.18)-(2.20). In canonical condensed notations this boundary value problem reads as

$$F_{ca}(\partial) G_{\text{DN}}^{ab}(y, y') = \delta_c^b \delta(y - y'), \quad (6.14)$$

$$G_{\text{DN}}^{ib}(y, y') \Big|_y = 0, \quad (6.15)$$

$$\vec{H}_a^A(\partial) G_{\text{DN}}^{ab}(y, y') \Big|_y = 0. \quad (6.16)$$

Thus, contrary to the purely Dirichlet ghost propagator the metric propagator G_{DN}^{ab} has Dirichlet conditions only for $i = (\alpha\beta, X)$ components. The rest of boundary conditions (6.16) belong to the generalized Neumann type. This is a consequence of the fact that lapse and shift functions in the path integral for $\Psi_B(g)$ are integrated out on the brane.

6.2. Brane effective action

Now substitute one-loop bulk wavefunction (6.12) into the path integral (5.2) and calculate it by the stationary-phase method. The stationary point of the overall tree-level phase satisfies the following equation

$$\frac{\partial}{\partial g^i} \left(S^B[G_0(g)] + S^b(g) + S_{\text{gb}}^b(g) \right) = \frac{\partial L_B}{\partial \dot{G}^i} \Big| + \frac{\partial S^b}{\partial g^i} - \frac{\partial \chi^\mu}{\partial g^i} c_{\mu\nu} \chi^\nu = 0, \quad (6.17)$$

because $S^B[G_0(g)]$ plays the role of the Hamilton-Jacobi function in the y -time canonical formalism, and its gradient in g^i yields the canonical momentum on the brane

$$p_i^0(G, \dot{G}) = \partial L_B / \partial \dot{G}^i,$$

$$\frac{\partial}{\partial g^i} S^B[G_0(g)] = \left. \frac{\partial L_B}{\partial \dot{G}^i} \right|_{G=G_0(g)}. \quad (6.18)$$

In essence (6.17) represents the generalized Israel matching condition relating the extrinsic curvature of the brane, $K^{\alpha\beta}(x) \sim \partial L_B / \partial \dot{G}_{\alpha\beta}(x)$, to the brane stress tensor $\partial S_b / \partial g^i = \delta S_b / \delta g_{\alpha\beta}(x)$ in the presence of the brane gauge-breaking term. To handle the latter, contract (6.17) with R_μ^i and take into account that in view of (3.18) and (3.27) the first term in the contraction vanishes on bulk shell,

$$R_\mu^i \frac{\partial L_B}{\partial \dot{G}^i} = - \left. \frac{\delta S^B}{\delta N^\mu} \right|_{G=G_0(g)} \equiv 0, \quad (6.19)$$

whereas the second term vanishes in view of d -dimensional covariance of the brane action (3.28), so that finally

$$R_\mu^i \frac{\partial \chi^\lambda}{\partial g^i} c_{\lambda\sigma} \chi^\sigma \equiv \mathbf{J}_\sigma^\lambda c_{\lambda\sigma} \chi^\sigma = 0, \quad (6.20)$$

whence $\chi^\mu = 0$ due to invertibility of $\mathbf{J}_\sigma^\lambda$ and $c_{\lambda\sigma}$. Therefore, we get the following system of equations for the stationary point g_0 in the d -dimensional brane gauge

$$\left(\left. \frac{\partial L_B}{\partial \dot{G}^i} \right| + \frac{\partial S^b}{\partial g^i} \right)_{g=g_0} = 0, \quad (6.21)$$

$$\chi^\mu(g_0) = 0. \quad (6.22)$$

To find the quadratic form of the tree-level action on the background of g_0 we would need the derivative of $G_0^a(g)$ with respect to g^i . This quantity satisfies the linearized version of the boundary-value problem (2.13)-(2.15). In condensed notations this problem for the bulk perturbation $\phi^a(\varphi)$ induced by the perturbation φ^i on the boundary reads as

$$\vec{S}_{ab}(\partial) \phi^b(y) = 0, \quad (6.23)$$

$$\vec{H}_b^A(\partial) \phi^b(y) = 0, \quad \phi^i| = \varphi^i. \quad (6.24)$$

To solve it notice that the Ab component of the Wronskian operator (6.14) is entirely determined by the gauge-breaking term of the full action

$$\vec{W}_{Ab}(\partial) = - \frac{\partial H^B}{\partial \dot{N}^A} c_{BD} \vec{H}_b^D(\partial) = a_{AD} \vec{H}_b^D(\partial), \quad (6.25)$$

$$a_{AD} \equiv a_A^B c_{BD}. \quad (6.26)$$

Therefore the problem (6.23)-(6.24) can be identically rewritten in the form

$$\vec{F}_{ab}(\partial) \phi^b(y) = 0, \quad (6.27)$$

$$\vec{W}_{Ab}(\partial) \phi^b| = 0, \quad \phi^i| = \varphi^i, \quad (6.28)$$

where the nondegenerate operator F_{ab} replaces the degenerate operator S_{ab} defined by (6.8). This is a problem with the (inhomogeneous) Dirichlet-Neumann boundary conditions. By using the Wronskian relation (2.26) its solution ϕ_{DN}^a can be presented in terms of the Dirichlet-Neumann Green's function of the above type

$$\phi_{\text{DN}}^a(y) = -G_{\text{DN}}^{ab}(y, y') \vec{W}_{bi}(\partial')|_{y'} \varphi^i \equiv -G_{\text{DN}}^{ab} \vec{W}_{bi}| \varphi^i, \quad (6.29)$$

whence

$$\frac{\partial G_0^a(g)}{\partial g^i} = -G_{\text{DN}}^{ab} \vec{W}_{bi}|. \quad (6.30)$$

Using (6.30) together with (6.18) and (3.24) one immediately obtains

$$\frac{\partial^2 S^{\text{B}}[G_0(g)]}{\partial g^i \partial g^k} = -\vec{W}_{ia} G_{\text{DN}}^{ab} \vec{W}_{bk}||, \quad (6.31)$$

where the double vertical bar denotes the restriction to the brane with respect to the both points of the two-point kernel, cf. Eq. (2.24) of Sect.2. This restriction, in particular, allows one to replace $\vec{W}_{ia}^S$ by $\vec{W}_{ia}$ in view of (6.16). As a result, the quadratic part of the full tree-level action on the background g_0 , $g = g_0 + \delta g$, takes the form

$$\frac{1}{2} \delta g^2 (S^{\text{B}}[G_0(g)] + S^{\text{b}}(g) + S_{\text{gb}}^{\text{b}}(g)) = \frac{1}{2} \delta g^i \mathbf{F}_{ik}^{\text{DN}} \delta g^k, \quad (6.32)$$

where $\mathbf{F}_{ik}^{\text{DN}}$ is the full brane-to-brane operator introduced in (2.23)

$$\mathbf{F}_{ik}^{\text{DN}} = -\vec{W}_{ia} G_{\text{DN}}^{ab} \vec{W}_{bk}|| + S_{ik}^{\text{b}} - \chi_i^\mu c_{\mu\nu} \chi_k^\nu, \quad (6.33)$$

with the gauge-fixed contribution of the brane action (2.27)

$$\kappa_{ik} = S_{ik}^{\text{b}} - \chi_i^\mu c_{\mu\nu} \chi_k^\nu, \quad S_{ik}^{\text{b}} \equiv \frac{\partial^2 S^{\text{b}}}{\partial g^i \partial g^k}, \quad \chi_i^\mu \equiv \frac{\partial \chi^\mu}{\partial g^i}. \quad (6.34)$$

Finally, substituting (6.12) into (5.2) and taking the Gaussian integral over δg^i in the vicinity of the stationary point g_0 one finds in the one-loop approximation

$$e^{i\Gamma(g_0)} = \frac{\mu_{\text{B}} \text{Det}_{\text{D}} Q_B^A}{(\text{Det}_{\text{DN}} F_{ab})^{1/2}} \frac{\mu_{\text{b}} \det \mathbf{J}_\nu^\mu}{(\det \mathbf{F}_{ik}^{\text{DN}})^{1/2}} \exp i(S^{\text{B}}[G] + S^{\text{b}}(g_0)) \Big|_{G=G_0(g_0)}. \quad (6.35)$$

The preexponential factor here confirms the decomposition property for the one-loop effective action (2.22) advocated in Sect.2 (modulo $\delta(0)$ -type terms generated by the local measure (4.15)).

7. Ward identities and gauge independence

It is important that the both bulk and brane preexponential factors in (6.35) are separately gauge independent. For the bulk prefactor this is a direct corollary of the gauge independence of $\Psi_B(g)$, (5.4). However, it is worth showing explicitly how this property works in virtue of the Ward identities for the bulk propagators.

Bulk Ward identities (for tree-level propagators) arise as follows [20]. Functional differentiation of (3.3) shows that on shell, that is on the background satisfying classical equations of motion, the functional matrix S_{ab} is degenerate because it has zero-eigenvalue eigenvectors – the gauge generators

$$R_A^a S_{ab}^B = -S_a^B \frac{\delta R_A^a}{\delta g^b} = 0 \quad (7.1)$$

As a consequence the operator F_{ab} satisfies the relation

$$R_A^a F_{ab} = -Q_A^B c_{BC} H_b^C \quad (7.2)$$

which can be functionally contracted with matrices of the gauge G^{bc} and ghost Q_D^{-1A} Green's functions. Integration by parts of the derivatives in the operators $R_A^a = R_A^a(\partial)$ and $Q_A^B = Q_A^B(\partial)$ does not produce additional surface terms in view of the boundary conditions for the propagators. Thus, one arrives at the identity relating the Dirichlet-Neumann Green's function of the metric operator to the Dirichlet Green's function of the Faddeev-Popov operator

$$c_{AB} \vec{H}_b^B(\partial) G_{\text{DN}}^{bc}(y, y') = -Q_A^{-1B}(y, y') \overleftarrow{R}_B^c(\partial_{y'}). \quad (7.3)$$

It is important that the Dirichlet conditions at y belonging to the boundary on the right hand side match with the Neumann conditions (6.16) on the left hand side.

Therefore, using the variational definitions of the functional determinants one has

$$\delta_H \ln \text{Det}_{\text{DN}} F_{ab} = \delta_H \overleftrightarrow{F}_{ab} G_{\text{DN}}^{ba} = -2 c_{AB} \vec{H}_b^B G_{\text{DN}}^{ba} \overleftarrow{H}_a^A, \quad (7.4)$$

$$\delta_H \ln \text{Det}_D Q_B^A = Q_A^{-1B} \delta_H \overleftarrow{Q}_B^A = Q_A^{-1B} \overleftarrow{R}_B^b \overleftarrow{H}_b^A, \quad (7.5)$$

so that in virtue of (7.3) the bulk part of the one-loop effective action is H -independent

$$\delta_H \left(-\frac{1}{2} \text{Tr}_{\text{DN}} \ln F + \text{Tr}_D \ln Q \right) = 0. \quad (7.6)$$

Bulk gauge obviously participates in the construction of the brane part of the effective action, but the latter turns out to be also independent of H^A . To show this, note first of all that in view of (6.16) $\vec{W}_{ia} G_{\text{DN}}^{ab} \overleftarrow{W}_{bk} = \vec{W}_{ia}^S G_{\text{DN}}^{ab} \overleftarrow{W}_{bk}^S$. This replacement

of the Wronskian operators $W(\partial)$ by their H -independent analogues $W^S(\partial)$, cf. Eq. (3.24), implies that

$$\delta_H \mathbf{F}_{ik}^{\text{DN}} = - \vec{W}_{ia}^S (\delta_H G_{\text{DN}}^{ab}) \overleftarrow{W}_{bk}^S \Big| \Big|. \quad (7.7)$$

The variation of the Green's function accounting for its boundary conditions is derived in Appendix C. Using (C.5) one has in view of (7.3)

$$\begin{aligned} \delta_H G_{\text{DN}}^{ab} &= -G_{\text{DN}}^{ac} \delta_H \overleftrightarrow{F}_{cd} G_{\text{DN}}^{db} = 2G_{\text{DN}}^{(ac} \overleftarrow{H}_c^A c_{AB} \delta \overrightarrow{H}_d^B G_{\text{DN}}^{db)} \\ &= -2 \overrightarrow{R}_A^a Q_B^{-1A} \delta \overrightarrow{H}_d^B G_{\text{DN}}^{db}, \end{aligned} \quad (7.8)$$

so that in virtue of the identity (3.26) the argument of the ghost propagator in

$$\begin{aligned} \delta_H \mathbf{F}_{ik}^{\text{DN}} &= 2 \vec{W}_{(ia}^S \overrightarrow{R}_A^a Q_B^{-1A} \delta H_d^B G_{\text{DN}}^{db} \overleftarrow{W}_{bk)} \Big| \Big|_{i,k} \\ &= -2 \frac{\partial \mathbf{H}_A}{\partial q^i} Q_B^{-1A} \delta H_d^B G_{\text{DN}}^{db} \overleftarrow{W}_{bk)} \Big| \Big|_{i,k} = 0 \end{aligned} \quad (7.9)$$

is not differentiated at the boundary because $\partial \mathbf{H}_A / \partial q^i$ is ultralocal in y , and the whole variation vanishes because of the Dirichlet boundary conditions for the ghost propagator, $(\partial \mathbf{H}_A / \partial q^i) Q_B^{-1A} \Big|_i = 0$. Finally,

$$\delta_H \left(-\frac{1}{2} \text{tr} \ln \mathbf{F}^{\text{DN}} + \text{tr} \ln \mathbf{J} \right) = 0. \quad (7.10)$$

Let us turn to the dependence of the effective action on the brane gauge χ^μ . As in usual gauge theories, the effective action is gauge-independent on shell, that is with the sources switched on the brane off.⁸ To see this in the one-loop order, note that the tree-level brane action is identically invariant with respect to d -dimensional (brane) diffeomorphisms

$$R_\mu^i \frac{\partial}{\partial g^i} (S^{\text{B}}[G_0(g)] + S^{\text{b}}(g)) \equiv 0. \quad (7.11)$$

The differentiation of this identity at $g = g_0$ (mass shell (6.21)) then gives

$$\begin{aligned} R_\mu^i \left(-\vec{W}_{ia} G_{\text{DN}}^{ab} \overleftarrow{W}_{bk} \Big| \Big| + S_{ik}^{\text{b}} \right) &= R_\mu^i \frac{\partial^2}{\partial g^i \partial g^k} (S^{\text{B}}[G_0(g)] + S^{\text{b}}(g)) \Big|_{g=g_0} \\ &= -\frac{\partial R_\mu^i}{\partial g^k} \frac{\partial}{\partial g^i} (S^{\text{B}}[G_0(g)] + S^{\text{b}}(g)) \Big|_{g=g_0} = 0. \end{aligned} \quad (7.12)$$

⁸By construction, there are no sources in the bulk, so that both $\Psi_{\text{B}}(g)$ and Γ are universally independent of the bulk gauge H , because the latter is not sensitive to the inclusion of boundary/brane sources. On the contrary, the brane gauge-fixing procedure is vulnerable for gauge non-invariant sources on the brane.

This means that the the brane-to-brane operator (6.33) is non-degenerate entirely due to its brane gauge-breaking term. Therefore, the following relation holds between the operators $\mathbf{F}_{ik}^{\text{DN}}$ and $\mathbf{J}_\mu^\nu$

$$R_\mu^i \mathbf{F}_{ik}^{\text{DN}} = -\mathbf{J}_\mu^\nu c_{\nu\alpha} \chi_k^\alpha. \quad (7.13)$$

Consequently, their Green's functions satisfy the brane Ward identity

$$\mathbf{J}_\nu^{-1\mu} R_\mu^i = -c_{\nu\alpha} \chi_k^\alpha \mathbf{G}^{ki}, \quad (7.14)$$

$$\mathbf{F}_{ik}^{\text{DN}} \mathbf{G}^{km} = \delta_i^m, \quad (7.15)$$

where $\mathbf{G}^{ki}$ is the brane-to-brane propagator. The relation (7.14) is a direct analogue of the bulk Ward identity (7.3), except that we don't have to care about integrations by parts in x , because the relevant surface terms are vanishing at the brane infinity $x^0 \rightarrow \pm\infty$ (or missing in Euclidean context in view of the closed nature of the boundary).

Thus, using the variations

$$\delta_\chi \mathbf{F}_{ik}^{\text{DN}} = -2\delta\chi_{(i}^\nu c_{\nu\alpha} \chi_{k)}^\alpha, \quad \delta_\chi \mathbf{J}_\mu^\nu = R_\mu^i \delta\chi_i^\nu \quad (7.16)$$

we immediately obtain in view of (7.14) the on-shell gauge independence of the brane effective action

$$\delta_\chi \left(-\frac{1}{2} \text{tr} \ln \mathbf{F}^{\text{DN}} + \text{tr} \ln \mathbf{J} \right) = \delta\chi_i^\nu c_{\nu\alpha} \chi_k^\alpha \mathbf{G}^{ki} + R_\mu^i \delta\chi_i^\nu \mathbf{J}_\nu^{-1\mu} = 0. \quad (7.17)$$

8. Reduction of the Dirichlet-Neumann problem to the Dirichlet type

If we would apply the stationary phase method directly to the path integral (4.11) without using the bulk wavefunction representation (5.2), then the one-loop contribution of the metric field would be determined by the functional determinant subject to Neumann boundary conditions for all metric components. So the decomposition of the integration procedure (5.1) exercises a half-way reduction to the Dirichlet problem, because the resulting boundary conditions are of a mixed type. As it was discussed in Introduction and in [1], the construction of the Dirichlet Green's function is much easier than the Neumann one, so actually no calculational advantages are gained when this reduction is incomplete. Therefore, it is worth making a further reduction from the Dirichlet-Neumann problem to the purely Dirichlet one.

For this purpose introduce the Gaussian path integral over metric perturbations $\delta G^a \equiv \phi^a = (\phi^i, \phi^A)$ with slightly more general than in (6.11) – inhomogeneous –

boundary conditions, $\phi^i| = \varphi^i$,

$$Z(\varphi^i) = \int_{\phi^i| = \varphi^i} D\phi \exp(-S[\phi]), \quad (8.1)$$

$$S[\phi] \equiv \frac{1}{2} \int dy \phi^a \overleftrightarrow{F}_{ab} \phi^b. \quad (8.2)$$

Its stationary point $\phi_{\text{DN}}^a(\varphi^i)$ is determined by the Dirichlet-Neumann problem with the inhomogeneous boundary conditions (6.27)-(6.28) and has the form (6.29). As shown in Appendix B, the gaussian integral itself equals

$$Z(\varphi^i) = (\text{Det } G_{\text{DN}})^{1/2} \exp(-S[\phi_{\text{DN}}(\varphi^i)]), \quad (8.3)$$

where the action at the stationary point is

$$S[\phi_{\text{DN}}(\varphi^i)] = -\frac{1}{2} \varphi^i \left(\overrightarrow{W}_{ia} G_{\text{DN}}^{ab} \overleftarrow{W}_{bk} \right) \varphi^k. \quad (8.4)$$

Alternatively it can be calculated by integrating first over the fields subject to Dirichlet boundary conditions for *all* ϕ^a with the integration over $\varphi^A \equiv \phi^A|$ reserved to the last

$$Z(\varphi^i) = \int \prod_A d\varphi^A \int_{\phi^a| = \varphi^a} D\phi \exp(-S[\phi]). \quad (8.5)$$

The inner integral

$$\int_{\phi^a| = \varphi^a} D\phi \exp(-S[\phi]) = (\text{Det } G_{\text{D}})^{1/2} \exp(-S[\phi_{\text{D}}(\varphi^a)]) \quad (8.6)$$

is determined by the Dirichlet problem for the field $\phi_{\text{D}}^a(\varphi)$, $F_{ab}\phi_{\text{D}}^b = 0$, $\phi_{\text{D}}^a| = \varphi^a$, having as a solution the following expression in terms of the Dirichlet Green's function G_{D} of the operator F_{ab}

$$\phi_{\text{D}}^a(\varphi) = -G_{\text{D}}^{ab} \overleftarrow{W}_{bc} \varphi^c|_c. \quad (8.7)$$

The action in (8.6) equals

$$S[\phi_{\text{D}}(\varphi^a)] = \frac{1}{2} \varphi^a \mathbf{D}_{ab} \varphi^b, \quad (8.8)$$

$$\mathbf{D}_{ab} \equiv -\overrightarrow{W}_{ac} G_{\text{D}}^{cd} \overleftarrow{W}_{db}|, \quad (8.9)$$

so that the gaussian integral over the boundary field φ^A in (8.5) is saturated by the saddle point φ_0^A of the above quadratic form in φ^A , $\varphi_0^A = -\mathbf{G}^{AB} \mathbf{D}_{Bi} \varphi^i$, where $\mathbf{D}_{Bi}$ is the Bi -block of the operator (8.9) and $\mathbf{G}^{AB}$ is the inverse of its AB -block

$$\mathbf{D}_{AB} \mathbf{G}^{BC} = \delta_A^C \quad (8.10)$$

(note that the operator (8.9) differs from the kernel of the action (8.4) by the type of the Green's function – Dirichlet vs Dirichlet-Neumann one).

Thus, integration over φ^A in (8.5) gives

$$Z(\varphi^i) = \left(\text{Det } G_D \right)^{1/2} \left(\det \mathbf{D}_{AB} \right)^{-1/2} \exp(-S[\phi_D(\varphi_0)]), \quad (8.11)$$

$$S[\phi_D(\varphi_0)] = \frac{1}{2} \varphi^i (\mathbf{D}_{ik} - \mathbf{D}_{iA} \mathbf{G}^{AB} \mathbf{D}_{Bk}) \varphi^k. \quad (8.12)$$

Comparison with (8.3)-(8.4) then gives the following two relations

$$-(\vec{W} G_{DN} \overleftarrow{W} ||)_{ik} = \mathbf{D}_{ik} - \mathbf{D}_{iA} \mathbf{G}^{AB} \mathbf{D}_{Bk}, \quad (8.13)$$

$$\left(\text{Det } G_{DN} \right)^{1/2} = \left(\text{Det } G_D \right)^{1/2} \left(\det \mathbf{D}_{AB} \right)^{-1/2}. \quad (8.14)$$

They allow one to completely reduce the algorithm (6.35) to that of the Dirichlet boundary conditions. In particular, the substitution of (8.13) into the expression for the brane-to-brane operator $\mathbf{F}_{ik}^{DN}$, (6.33), expresses its functional determinant as

$$\det \mathbf{F}_{ik}^{DN} = \det \left((\mathbf{D}_{ik} + \kappa_{ik}) - \mathbf{D}_{iA} \mathbf{G}^{AB} \mathbf{D}_{Bk} \right) = \det \mathbf{F}_{ab}^D \left(\det \mathbf{D}_{AB} \right)^{-1}, \quad (8.15)$$

where $\mathbf{F}_{ab}^D$ is the brane-to-brane operator of another type – it acts on the space of all metric perturbations on the brane $\delta G^a| = (\delta g^i, \delta N^A)|$, $a = (i, A)$, and is built of the Dirichlet Green's function rather than the mixed-type one

$$\mathbf{F}_{ab}^D = \begin{bmatrix} \mathbf{D}_{ik} + \kappa_{ik} & \mathbf{D}_{iB} \\ \mathbf{D}_{Ak} & \mathbf{D}_{AB} \end{bmatrix} = -(\vec{W} G_D \overleftarrow{W} ||)_{ab} + \kappa_{ik} \delta_a^i \delta_b^k. \quad (8.16)$$

Its ik -block is given by Eq.(2.29) in Sect.2.

In view of the relation (8.15) the one-loop contribution of the bulk and brane gravitons in (6.35) can be decomposed into the product of other two bulk and brane factors which are entirely based on the Dirichlet-type objects

$$\left(\text{Det}_{DN} F \right)^{-1/2} \left(\det \mathbf{F}_{ik}^{DN} \right)^{-1/2} = \left(\text{Det}_D F \right)^{-1/2} \left(\det \mathbf{F}_{ik}^D \right)^{-1/2}. \quad (8.17)$$

In fact, this decomposition literally repeats the Neumann-Dirichlet reduction suggested in [1]. The corresponding decomposition of the one-loop effective action has the form (2.28) presented in Sect.2. From calculation viewpoint it is simpler than (2.22), but it destroys manifest gauge independence of the combined bulk-brane diagrammatic technique. Whereas in (2.22) both bulk and brane parts are separately gauge-independent, no such property holds for (2.28) – only the sum of bulk and brane terms is independent of the bulk gauge conditions.

9. Conclusions

Thus we have an exhaustive formulation of the Feynman-DeWitt-Faddeev-Popov gauge-fixing procedure in gravitational systems with branes or boundaries. This procedure incorporates a special choice of gauge conditions in the bulk and on the brane, which separately fix the bulk and brane diffeomorphisms. Also it establishes the boundary conditions for the corresponding ghost factors and allows one to construct a special bulk wavefunction representation of the brane effective action (5.2)-(5.3). The bulk wavefunction satisfies the generalized Wheeler-DeWitt equations with respect to the induced metric of the brane (5.7). They might, perhaps, serve as a basis for non-perturbative methods in brane models, alternative to semiclassical expansion [14].

We derived the boundary conditions for propagators of the Feynman diagrammatic technique in brane models and explicitly built the brane effective action in the one-loop approximation. Similarly to non-gauge (quantized matter) models with branes, considered in [1], this one-loop action can be decomposed into the sum of bulk and brane contributions both in the graviton and the ghost sectors (2.28). The bulk contribution is represented by the usual functional determinant of the propagator in the $(d + 1)$ -dimensional bulk, subject to Dirichlet boundary conditions on the brane. The brane contribution is given by a similar determinant of the brane-to-brane operator (2.29) in the surface d -dimensional theory. In the graviton sector this operator has a nonlocal (pseudodifferential) nature, whereas in the ghost sector for local gauges this is always a local Faddeev-Popov operator associated with gauging away the d -dimensional surface diffeomorphisms. This property follows from the generic bulk-brane factorization of a gauge-fixing factor (2.5) in the path integral for gravitational brane models, which holds beyond loop expansion.

Linear algebra manipulations and Gaussian integrations which underly the above results look innocent at the calculational level adopted in this paper. The obtained algorithms are, however, marred by ultraviolet divergences and should be regulated by some covariant technique. The efficiency and correctness of these calculations was, nevertheless, confirmed within the dimensional regularization in [1]. In particular, correct expressions for surface Schwinger-DeWitt coefficients were obtained in [1] and, thus, guaranteed correct renormalization of logarithmic divergences in simplest brane models.

Such calculations should also apply in gravitational systems considered above, and this is a subject of forthcoming papers. We plan to make a synthesis of the Neumann-Dirichlet reduction with a systematic curvature expansion method in brane models in order to alleviate their formalism to the level of universality of the Schwinger-DeWitt technique [23, 24]. This is important for various applications in quantum gravity and

cosmology, including quantum consistency of brane models [6], the conformal anomaly mechanism of the dark energy, that might be facilitated within the brane concept of extra dimensions [7], and the others.

Appendix A The local measure in brane models

Let the bulk and brane terms of the action both have kinetic terms quadratic in velocities with coefficients $a(X)$ and $b(x)$. These terms can be rewritten as one integral over the bulk

$$\begin{aligned} \int_{\mathbf{B}} dX a(X) (\partial_t \phi(X))^2 + \int_{\mathbf{b}} dx b(x) (\partial_t \varphi(x))^2 \\ = \int_{\mathbf{B}} dX dX' \mathcal{G}(X, X') \partial_t \phi(X) \partial_t \phi(X') \end{aligned} \quad (\text{A.1})$$

with the overall coefficient of the quadratic form in field velocities

$$\mathcal{G}(X, X') = \delta(x, x') [a(X) \delta(y - y') + b(x) \delta(y) \delta(y')] . \quad (\text{A.2})$$

Here $\phi(X)$ and $\varphi(x) = \phi(x, 0) = \phi(X)|$ denote the bulk field and its boundary value at the brane located at $y = 0$.

The full local canonical measure reads in terms of the functional determinant of this matrix as

$$\mu[a(X), b(x)] \equiv \left(\text{Det } \mathcal{G}(X, X') \right)^{1/2} = \mu_B[a(X)] \mu_b[b(x)] \quad (\text{A.3})$$

and as we show below factorizes into the product of the corresponding bulk $\mu_B[a(X)]$ and brane $\mu_b[b(x)]$ measures given by

$$\mu_B[a(X)] = \left(\frac{\text{Det}_{(d+1)} a(X) \delta(X, X')}{\text{Det}_{(d)} a(x, 0) \delta(x, x')} \right)^{1/2} \quad (\text{A.4})$$

and

$$\mu_b[b(x)] = \left(\text{Det}_{(d)} b(x) \delta(x, x') \right)^{1/2} \quad (\text{A.5})$$

This factorization follows from the expression for the inverse of the functional matrix (A.2)

$$\mathcal{G}^{-1}(X, X') = \delta(x, x') \frac{1}{a(X)} \left[\delta(y - y') - \frac{b(x) \delta(y) \delta(y')}{a(x, 0) + \delta(0) b(x)} \right] . \quad (\text{A.6})$$

When substituted into the variational equation for the logarithm of the functional determinant in (A.3) this expression yields the decomposition into the following sum of two terms

$$\begin{aligned}\delta \ln \text{Det } \mathcal{G}(X, X') &= \int dX dX' \delta \mathcal{G}(X, X') \mathcal{G}^{-1}(X', X) \\ &= \delta \left[\delta^{(d+1)}(0) \int_{\mathbf{B}} dX \ln a(X) + \delta^{(d)}(0) \int_{\mathbf{b}} dx \ln \left(1 + \delta(0) \frac{b(x)}{a(x, 0)} \right) \right].\end{aligned}\quad (\text{A.7})$$

With the obvious limit⁹ $\delta(0) \rightarrow \infty$ this sum gives the factorization of the functional determinant

$$\text{Det } \mathcal{G}(X, X') = \frac{\text{Det}_{(d+1)} a(X) \delta(X, X')}{\text{Det}_{(d)} a(x, 0) \delta(x, x')} \text{Det}_{(d)} b(x) \delta(x, x'), \quad (\text{A.8})$$

which implies the factorization (A.3)-(A.5). In its turn this confirms the factorization of the local measure in the brane model (4.15)-(4.17). The division by $\text{Det}_{(d)} a(x, 0) \delta(x, x')$ in the bulk measure (A.4) corresponds to the absence of local factors belonging to the brane in the product of (4.16). In the notations of this Appendix this means the following chain of relations

$$\begin{aligned}\prod_{X, X \notin \mathbf{b}} a(X) &= \prod_{y > 0, x} a(x, y) = \prod_x \frac{1}{a(x, 0)} \prod_{y \geq 0} a(x, y) \\ &= \prod_x \frac{1}{a(x, 0)} \exp \left(\delta(0) \int_{y \geq 0} dy \ln a(x, y) \right) = \frac{\text{Det}_{(d+1)} a(X) \delta(X, X')}{\text{Det}_{(d)} a(x, 0) \delta(x, x')}.\end{aligned}\quad (\text{A.9})$$

Note that this division is responsible for zero density weight of the bulk wavefunction $\Psi_{\mathbf{B}}(g)$, defined by the path integral (5.3) (see footnote 7 in Sect.5).

Appendix B The Gaussian functional integral with mixed boundary conditions

Feynman's calculation [22] of the gaussian functional integral with the action (6.11) for the case when only a part of integration fields is fixed at the boundary,

$$\delta G^a \equiv \phi^a = (\phi^i, \phi^A), \quad \phi^i| = 0, \quad (\text{B.1})$$

⁹Any regularization implies this limit except the dimensional regularization in which $\delta(0) = 0$, but in the dimensional regularization the contribution of the local measure is identically zero.

can be based on the integral

$$Z[F, J] = \int_{\phi^i|_{=0}} D\phi \exp(-S[\phi, J]), \quad (\text{B.2})$$

$$S[\phi, J] = \int dy \left(\frac{1}{2} \phi^a(y) \overleftrightarrow{F}_{ab}(\partial) \phi^b(y) + J_a(y) \phi^a(y) \right), \quad (\text{B.3})$$

where the action is modified by the source term in the bulk. To find the dependence of (B.2) on J_a consider the stationary point $\phi_{\text{DN}} = \phi_{\text{DN}}[J]$ of this action with respect to variations of ϕ^a satisfying (B.1).

$$\delta S[\phi, J] = \int dy \delta\phi^a (\overrightarrow{F}\phi + J)_a + \delta\phi^A (\overrightarrow{W}\phi)_A \Big| = 0 \quad (\text{B.4})$$

Thus, ϕ_{DN} satisfies the problem with mixed Dirichlet-Neumann boundary conditions

$$\overrightarrow{F}_{ab} \phi_{\text{DN}}^b + J_a = 0, \quad (\text{B.5})$$

$$\phi_{\text{DN}}^i|_{=0}, \quad \overrightarrow{W}_{Aa} \phi_{\text{DN}}^a|_{=0}, \quad (\text{B.6})$$

and therefore can be represented in terms of the corresponding Green's function of Eqs.(6.14)-(6.16)

$$\phi_{\text{DN}}^a[J] = G_{\text{DN}}^{ab} J_b \equiv \int dy' G_{\text{DN}}^{ab}(y, y') J_b(y') \quad (\text{B.7})$$

Now make the shift of the integration variable in (B.2) by ϕ_{DN} , $\phi = \phi_{\text{DN}} + \Delta$. Under this replacement the action decomposes in the part $S[\Delta, 0]$ quadratic in Δ and the part independent of Δ . Linear in Δ term is absent (both in the bulk and on the boundary) in view of the stationarity of the action at ϕ_{DN} , so that

$$S[\phi, J] = S[\Delta, 0] + S[\phi_{\text{DN}}, J], \quad (\text{B.8})$$

$$S[\phi_{\text{DN}}, J] = \frac{1}{2} J_a G_{\text{DN}}^{ab} J_b. \quad (\text{B.9})$$

Therefore

$$Z[F, J] = Z[F, 0] \exp(-S[\phi_{\text{DN}}, J]). \quad (\text{B.10})$$

To find the prefactor, consider the variation of the integral (B.2) at $J = 0$ with respect to the operator F_{ab} and make the following set of obvious identical transformations using the above equations (B.9) and (B.10)

$$\begin{aligned} \delta_F Z[F, 0] &= - \int D\phi \left(\frac{1}{2} \phi^a \overleftrightarrow{\delta F}_{ab} \phi^b \right) \exp(-S[\phi, 0]) \\ &= - \int D\phi \left(\frac{1}{2} \frac{\delta}{\delta J_a} \overleftrightarrow{\delta F}_{ab} \frac{\delta}{\delta J_b} \right) Z[F, J] \Big|_{J=0} \\ &= -\frac{1}{2} \overleftrightarrow{\delta F}_{ab} G_{\text{DN}}^{ba} Z[F, 0]. \end{aligned} \quad (\text{B.11})$$

Here $\overleftrightarrow{\delta F}_{ab}$ means arbitrary variations of the coefficients of the operator, and the double arrow implies symmetric action of two first-order derivatives of $F_{ab}(\partial)$ on both arguments of G_{DN}^{ba} similar to Eq.(B.3).

Therefore, one gets

$$\delta_F \ln Z[F, 0] = -\frac{1}{2} \overleftrightarrow{\delta F}_{ab} G_{\text{DN}}^{ba} = -\frac{1}{2} \delta \ln \text{Det}_{\text{DN}} F \equiv \frac{1}{2} \delta \ln \text{Det} G_{\text{DN}}. \quad (\text{B.12})$$

This justifies the variational definition (6.13) of the Dirichlet-Neumann functional determinant of F_{ab} originating from the Gaussian integration with mixed boundary conditions.

Appendix C The variation of G_{DN}^{ab}

In view of (6.25) $\vec{H}_a^A = a^{-1AC} \vec{W}_{Cb}$, and the boundary conditions in the problem (6.14)-(6.16) for G_{DN}^{ab} can be rewritten in terms of the Wronskian operator $\vec{W}_{Cb}$. Then the variation of this problem gives the set of inhomogeneous equations for $\delta G_{\text{DN}}^{ab}$

$$\begin{aligned} \vec{F}_{ab} \delta G_{\text{DN}}^{bc} &= -\delta \vec{F}_{ab} G_{\text{DN}}^{bc}, \\ \vec{W}_{Ab} \delta G_{\text{DN}}^{bc} \Big|_A &= -\delta \vec{W}_{Ab} G_{\text{DN}}^{bc} \Big|_A, \\ \delta G_{\text{DN}}^{ic} \Big|_i &= 0 \end{aligned} \quad (\text{C.1})$$

This set of equations forms the boundary value problem, analogous to (6.27)-(6.29), but with the inhomogeneous subset of Neumann boundary conditions

$$\begin{aligned} \vec{F} \phi &= J, \\ (\vec{W} \phi)_A &= w_A, \quad \phi^i \Big|_i = 0. \end{aligned} \quad (\text{C.2})$$

It has a solution

$$\begin{aligned} \phi^a(X) &= \int_{\mathbf{B}} dX' G_{\text{DN}}^{ab}(X, X') J_b(X') + \int_{\mathbf{b}} dx G_{\text{DN}}^{aA}(X, e(x)) w_A(x) \\ &\equiv G_{\text{DN}}^{ab} J_b + G_{\text{DN}}^{aA} w_A \Big|_A \end{aligned} \quad (\text{C.3})$$

which, when applied to (C.1), gives

$$\delta G_{\text{DN}}^{ab} = -G_{\text{DN}}^{ac} \delta \vec{F}_{cd} G_{\text{DN}}^{db} - G_{\text{DN}}^{aA} \delta \vec{W}_{Ad} G_{\text{DN}}^{db} \Big|_A \quad (\text{C.4})$$

Therefore, in view of the relation (6.10), the variation of the Dirichlet-Neumann Green's function is given by the equation

$$\delta G_{\text{DN}}^{ab} = -G_{\text{DN}}^{ac} \overleftrightarrow{\delta F}_{cd} G_{\text{DN}}^{db}, \quad (\text{C.5})$$

used in Eq.(7.8).

Acknowledgements

The author is grateful for hospitality of the Physics Department of the University of Bologna and the Theoretical Physics Institute of the University of Cologne. This work was supported by the Russian Foundation for Basic Research under the grant No 05-01-00996 and the grant LSS-4401.2006.2.

References

- [1] A.O.Barvinsky and D.V.Nesterov, Phys. Rev. **D73** (2006) 066012, hep-th/0512291.
- [2] J.Garriga and T.Tanaka Phys. Rev. Lett. **84** (2000) 2778, hep-th/9911055.
- [3] L.Pilo, R.Rattazzi and A.Zaffaroni, JHEP **0311** (2000) 056, hep-th/0004028.
- [4] S.L.Dubovsky and V.A.Rubakov, Phys. Rev. **D 67** (2003) 104014, hep-th/0212222; S.L.Dubovsky and M.V.Libanov, JHEP **0311** (2003) 038, hep-th/0309131.
- [5] M.A.Luty, M.Porrati and R.Rattazzi, JHEP **0309** (2003) 029, hep-th/0303116.
- [6] A.Nicolis and R.Rattazzi, JHEP **06** (2004) 059, hep-th/0404159.
- [7] A.O.Barvinsky and A.Yu.Kamenshchik, *Cosmological landscape from nothing: some like it hot*, hep-th/0605132.
- [8] L.Randall and S.Sundrum, Phys. Rev. Lett. **83** (1999) 4690, hep-th/9906064.
- [9] G.R.Dvali, G.Gabadadze and M.Porrati, Phys. Lett. **B485** (2000) 208, hep-th/0005016.
- [10] T.Tanaka, Phys. Rev. **D69** (2004) 024001, gr-qc/0305031.
- [11] M.Kolanovic, M.Porrati and J.W.Rombouts, Phys. Rev. **D68** (2003) 064018; M.Porrati and J.W.Rombouts, Phys. Rev. **D69** (2004) 122003, hep-th/0401211.
- [12] B.S.DeWitt, *Dynamical Theory of Groups and Fields* (Gordon and Breach, New York, 1965); Phys.Rev. **162** (1967) 1195.
- [13] A.O.Barvinsky and D.V.Nesterov, Nucl. Phys. **B654** (2003) 225, hep-th/0210005.
- [14] A.O.Barvinsky, *The Gospel according to DeWitt revisited: quantum effective action in braneworld models*, hep-th/0504205.

- [15] E.S.Fradkin and G.A.Vilkovisky, Phys.Lett. **B55** (1975) 224, CERN Report TH-2332 (1977).
- [16] H.Leutwyler, Phys. Rev. **134** (1964) B1155.
- [17] J.B.Hartle and S.W.Hawking, Phys.Rev. **D28** (1983) 2960.
- [18] A.O.Barvinsky and V.N.Ponomariov, Phys.Lett. **B167** (1986) 289; A.O.Barvinsky, Phys.Lett. **B195** (1987) 344; Phys. Rep. **230** (1993) 237.
- [19] J.J.Halliwell and J.B.Hartle, Phys.Rev. **D43** (1991) 1170.
- [20] A.O.Barvinsky, Nucl. Phys. **B 520** (1998) 533.
- [21] A.O.Barvinsky, Phys. Lett. **B241** (1990) 201; *Geometry of the Dirac and reduced phase space quantization of constrained systems*, gr-qc/9612003; A.O.Barvinsky and V.Krykhtin, Class.Quantum Grav. **10** (1993) 1957.
- [22] R.P.Feynman, Phys.Rev. **84** (1951) 108.
- [23] A.O. Barvinsky and G.A. Vilkovisky, Phys. Rep. 119 (1985) 1.
- [24] A.O.Barvinsky and G.A.Vilkovisky, Nucl. Phys. **B 282** (1987) 163, Nucl. Phys. **B333** (1990) 471; A.O.Barvinsky, Yu.V.Gusev, G.A.Vilkovisky and V.V.Zhytnikov, J. Math. Phys. **35** (1994) 3525; J. Math. Phys. **35** (1994) 3543.